

EXPLICIT EVALUATION OF CERTAIN EXPONENTIAL SUMS OF QUADRATIC FUNCTIONS OVER $\mathbb{F}_{p^n}$, p ODD

SANDRA DRAPER AND XIANG-DONG HOU

ABSTRACT. Let p be an odd prime and let $f(x) = \sum_{i=1}^k a_i x^{p^{\alpha_i}+1} \in \mathbb{F}_{p^n}[x]$, where $0 \leq \alpha_1 < \cdots < \alpha_k$. We consider the exponential sum $S(f, n) = \sum_{x \in \mathbb{F}_{p^n}} e_n(f(x))$, where $e_n(y) = e^{2\pi i \text{Tr}_n(y)/p}$, $y \in \mathbb{F}_{p^n}$, $\text{Tr}_n = \text{Tr}_{\mathbb{F}_{p^n}/\mathbb{F}_p}$. There is an effective way to compute the nullity of the quadratic form $\text{Tr}_{mn}(f(x))$ for all integer $m > 0$. Assuming that all such nullities are known, we find relative formulas for $S(f, mn)$ in terms of $S(f, n)$ when $\nu_p(m) \leq \min\{\nu_p(\alpha_i) : 1 \leq i \leq k\}$, where ν_p is the p -adic order. We also find an explicit formula for $S(f, n)$ when $\nu_2(\alpha_1) = \cdots = \nu_2(\alpha_k) < \nu_2(n)$. These results generalize those by Carlitz and by Baumert and McEliece. Parallel results with $p = 2$ were obtained in a previous paper by the second author.

1. INTRODUCTION

Let p be a prime and let $\mathbb{F}_{p^n}$ be the finite fields with p^n elements. Denote the trace from $\mathbb{F}_{p^n}$ to $\mathbb{F}_{p^m}$ by $\text{Tr}_{n/m}$; the trace $\text{Tr}_{n/1}$ is also denoted by Tr_n . Let $e_n(y) = e^{2\pi i \text{Tr}_n(y)/p}$, $y \in \mathbb{F}_{p^n}$. Carlitz considered the sum $\sum_{x \in \mathbb{F}_{p^n}} e_n(ax^{p+1} + bx)$, where $a, b \in \mathbb{F}_{p^n}$. He gave explicit evaluations of this sum for $p = 2$ in [2] and for odd p in [3]. The evaluation of the sum $\sum_{x \in \mathbb{F}_{p^n}} e_n(ax^{p^\alpha+1})$, where $a \in \mathbb{F}_{p^n}$ and $\alpha \geq 0$, was also implied by the results of Baumert and McEliece [1]. Also see [5].

More generally, we let

$$(1.1) \quad f(x) = \sum_{i=1}^k a_i x^{p^{\alpha_i}+1} \in \mathbb{F}_{p^n}[x],$$

where $0 \leq \alpha_1 < \cdots < \alpha_k$, and consider the sum

$$S(f, n) = \sum_{x \in \mathbb{F}_{p^n}} e_n(f(x)).$$

Note that $S(f, n)$ is an exponential sum of a quadratic form on $\mathbb{F}_p^n$. In fact, if we identify $\mathbb{F}_{p^n}$ with $\mathbb{F}_p^n$, then $\text{Tr}_n(f(x))$ is a quadratic form in the coordinates of $\mathbb{F}_p^n$. Conversely, every quadratic form on $\mathbb{F}_{p^n}$ (identified with $\mathbb{F}_p^n$) can be written as $\text{Tr}_n(f(x))$ for some polynomial of the form (1.1). The sum $S(f, n)$ is completely determined by the canonical form of the quadratic form $\text{Tr}_n(f(x))$. The converse is true except when $p = 2$ and $S(f, n) = 0$. In general, there is no direct way to identify the canonical form of $\text{Tr}_n(f(x))$ using $f(x)$; an indirect way is through the computation of $S(f, n)$.

1991 *Mathematics Subject Classification.* 11T23.

Key words and phrases. Exponential sum, Gauss sum, law of quadratic reciprocity, quadratic form.

We can also consider a seemingly more general sum

$$S(f + bx, n) = \sum_{x \in \mathbb{F}_{p^n}} e_n(f(x) + bx),$$

where $b \in \mathbb{F}_{p^n}$. However, when p is odd, $S(f + bx, n)$ follows easily from $S(f, n)$, as we will see in section 3. When $p = 2$, we can write $S(f + bx, n) = S(f + cx^2, n)$, where $b = c^2$ and $f + cx^2$ is of the form (1.1).

In general, the sum $S(f, n)$ is not explicitly known. In a recent paper [6] by the second author, Carlitz's result on the sum $\sum_{x \in \mathbb{F}_{2^n}} e_n(ax^3 + bx)$ was extended to certain polynomials of the form (1.1) in $\mathbb{F}_{2^n}[x]$. In the present paper, we will establish parallel results for odd primes p .

When p is odd, the sum $S(f, n)$ is expressed in terms of the nullity and type (definitions in the next section) of the quadratic form $\text{Tr}_n(f(x))$. The nullity of $\text{Tr}_n(f(x))$ is rather easy to compute, as we will see in the next section. For most of the paper, we will treat the nullity of $\text{Tr}_n(f)$ as known and focus on the determination of the type of $\text{Tr}_n(f(x))$.

We fix a polynomial $f(x) \in \mathbb{F}_{p^n}[x]$ of the form (1.1) and consider the sum $S(f, mn)$ as a function of $m \in \mathbb{Z}^+$. We try to evaluate the type of $\text{Tr}_{mn}(f(x))$ in terms of that of $\text{Tr}_n(f(x))$; the result gives $S(f, mn)$ in terms of $S(f, n)$. For this purpose, it suffices to assume that m is a prime, say q . The three cases, (i) $q \neq p, 2$, (ii) $q = 2$, and (iii) $q = p$, require different methods. In the first two cases, we obtain explicit formulas for $S(f, qn)$ in terms of $S(f, n)$. When $q = p$, we are able to express $S(f, pn)$ in terms of $S(f, n)$ only under a very restrictive condition that $\nu_p(n) < \min\{\nu_p(\alpha_i) : 1 \leq i \leq k\}$, where ν_p is the p -adic order function and $\alpha_1, \dots, \alpha_k$ are given in (1.1).

The results described above are “relative” formulas of $S(f, mn)$ in terms of $S(f, n)$. Explicit formulas for $S(f, n)$ itself would be more desirable. A special situation which allows explicit evaluation of $S(f, n)$ is when $\nu_2(\alpha) = \dots = \nu_2(\alpha_k) < \nu_2(n)$. A similar result for $p = 2$ has been proved in [6].

The paper is organized as follows. In section 2, we express $S(f, n)$ in terms of the nullity and type of $\text{Tr}_n(f)$. We also describe a method for computing the nullity of $\text{Tr}_n(f)$. Section 3 provides a formula for $S(f + bx, n)$ in terms of $S(f, n)$. In sections 4 – 6, we derive relative formulas for $S(f, mn)$ in terms of $S(f, n)$ in three cases respectively: (i) $m = q^s$, where q is a prime and $q \neq p, 2$; (ii) $m = 2^s$; (iii) $m = p^s$, where $s \leq \min\{\nu_p(\alpha_i) : 1 \leq i \leq k\} - \nu_p(n)$. In section 7, we give an explicit formula for $S(f, n)$ under the assumption that $\nu_2(\alpha_1) = \dots = \nu_2(\alpha_k) < \nu_2(n)$. As a special case of this result, we have an explicit formula for the sum $S(ax^{p^a+1})$. The two tables in the appendix contain the nullities of $\text{Tr}_n(f)$ with $p^n = 3$, $\alpha_k \leq 4$ and with $p^n = 5$, and $\alpha_k \leq 3$.

Throughout the paper, p is always an odd prime.

2. THE NULLITY AND TYPE

It is well known (cf. [8, Theorem 6.9]) that every quadratic form $F(x_1, \dots, x_n)$ on $\mathbb{F}_p^n$ is congruent to a canonical form

$$(2.1) \quad x_1^2 + \dots + x_{r-1}^2 + dx_r^2,$$

where $r \geq 0$ and $d \in \mathbb{F}_p^*$. (Two quadratic forms $F(x_1, \dots, x_n)$ and $G(x_1, \dots, x_n)$ on $\mathbb{F}_p^n$ are congruent if $G(x_1, \dots, x_n) = F((x_1, \dots, x_n)A)$ for some $A \in \text{GL}(n, \mathbb{F}_p)$.) In

(2.1), the integer r is called the rank of F and $n - r$ the nullity of F . Let η be the quadratic character of $\mathbb{F}_p$. Then $\eta(d) \in \{\pm 1\}$ is called the type of F . The image of d in the quotient group $\mathbb{F}_p^*/(\mathbb{F}_p^*)^2$, where $(\mathbb{F}_p^*)^2 = \{a^2 : a \in \mathbb{F}_p^*\}$, is called the discriminant of F . (If $F = 0$, we define the type of F to 1 and the discriminant of F to be $1(\mathbb{F}_p^*)^2$.) The nullity of F is the dimension of the $\mathbb{F}_p$ -space

$$\{z \in \mathbb{F}_p^n : F(x+z) - F(x) = \text{constant for all } x \in \mathbb{F}_p^n\}.$$

The quadratic form F is uniquely determined, up to congruence, by its rank and type.

Let g_p denote the Gauss quadratic sum on $\mathbb{F}_p$. Recall that

$$g_p = \sum_{x \in \mathbb{F}_p} \zeta_p^{x^2} = i^{\frac{1}{4}(p-1)^2} p^{\frac{1}{2}},$$

where $\zeta_p = e^{2\pi i/p}$.

Lemma 2.1. *Let F be a quadratic form on $\mathbb{F}_p^n$ of type t and rank r . Then*

$$\sum_{(x_1, \dots, x_n) \in \mathbb{F}_p^n} \zeta_p^{F(x_1, \dots, x_n)} = t g_p^r p^{n-r}.$$

Proof. We may assume $F(x_1, \dots, x_n) = x_1^2 + \dots + x_{r-1}^2 + dx_r^2$, where $\eta(d) = t$. Then

$$\sum_{(x_1, \dots, x_n) \in \mathbb{F}_p^n} \zeta_p^{F(x_1, \dots, x_n)} = p^{n-r} \left(\sum_{x \in \mathbb{F}_p} \zeta_p^{x^2} \right)^{r-1} \left(\sum_{x \in \mathbb{F}_p} \zeta_p^{dx^2} \right) = t g_p^r p^{n-r}.$$

□

Now let

$$(2.2) \quad f(x) = \sum_{i=1}^k a_i x^{p^{\alpha_i} + 1} \in \mathbb{F}_{p^n}[x],$$

where $0 \leq \alpha_1 < \dots < \alpha_k = \alpha$ and $a_k \neq 0$. Let $l_n(f)$ and $t_n(f)$ denote the nullity and type of $\text{Tr}_n(f)$ respectively. By Lemma 2.1,

$$(2.3) \quad S(f, n) = t_n(f) g_p^{n-l_n(f)} p^{l_n(f)}.$$

The nullity $l_n(f)$ is the dimension of the $\mathbb{F}_p$ -space

$$\{z \in \mathbb{F}_{p^n} : \text{Tr}_n(f(x+z) - f(x)) = \text{constant for all } x \in \mathbb{F}_{p^n}\}.$$

Note that

$$\begin{aligned}
& \text{Tr}_n(f(x+z) - f(x)) \\
&= \text{Tr}_n \left[\sum_{i=1}^k a_i ((x+z)^{p^{\alpha_i}+1} - x^{p^{\alpha_i}+1}) \right] \\
&= \text{Tr}_n \left[\sum_{i=1}^k a_i (x^{p^{\alpha_i}} z + x z^{p^{\alpha_i}} + z^{p^{\alpha_i}+1}) \right] \\
(2.4) \quad &= \text{Tr}_n(f(z)) + \text{Tr}_n \left[x \sum_{i=1}^k (a_i z^{p^{\alpha_i}} + a_i^{p^{-\alpha_i}} z^{p^{-\alpha_i}}) \right] \\
&= \text{Tr}_n(f(z)) + \text{Tr}_n \left[x^{p^\alpha} \sum_{i=1}^k (a_i^{p^\alpha} z^{p^{\alpha+\alpha_i}} + a_i^{p^{\alpha-\alpha_i}} z^{p^{\alpha-\alpha_i}}) \right] \\
&= \text{Tr}_n(f(z)) + \text{Tr}_n[x^{p^\alpha} f^*(z)],
\end{aligned}$$

where $p^{-\alpha_i}$ is a positive integer such that $p^{-\alpha_i} \cdot p^{\alpha_i} \equiv 1 \pmod{p^n - 1}$ and

$$(2.5) \quad f^*(z) = \sum_{i=1}^k (a_i^{p^\alpha} z^{p^{\alpha+\alpha_i}} + a_i^{p^{\alpha-\alpha_i}} z^{p^{\alpha-\alpha_i}}).$$

Therefore, $\text{Tr}_n(f(x+z) - f(x)) = \text{constant}$ for all $x \in \mathbb{F}_{p^n}$ if and only if z is a root of f^* . f^* is a p -polynomial over $\mathbb{F}_{p^n}$ without repeated roots. Thus,

$$\begin{aligned}
(2.6) \quad l_n(f) &= \dim_{\mathbb{F}_p} \{z \in \mathbb{F}_{p^n} : f^*(z) = 0\} \\
&= \log_p |\{z \in \mathbb{F}_{p^n} : f^*(z) = 0\}| \\
&= \log_p \deg(f^*, x^{p^n} - x).
\end{aligned}$$

Let

$$(2.7) \quad s = \min\{m : n \mid m, l_m(f) = 2\alpha\}.$$

Since $\deg f^* = p^{2\alpha}$, $\mathbb{F}_{p^s}$ is the splitting field of f^* over $\mathbb{F}_{p^n}$. It is obvious that

$$(2.8) \quad l_m(f) = l_{(m,s)}(f) \quad \text{for all } 0 < m \equiv 0 \pmod{n}.$$

The nullity $l_m(f)$ for all $0 < m \equiv 0 \pmod{n}$ can be computed using the following algorithm: First use (2.6) to compute $l_{in}(f)$ for $i = 1, 2, \dots$ until $l_s(f) = 2\alpha$. Then $\mathbb{F}_{p^s}$ is the splitting field of f^* and $l_m(f)$ is given by (2.8).

Example 2.2. Let

$$f(x) = x^{5^0+1} + 2x^{5^1+1} + 3x^{5^2+1} + 4x^{5^3+1} + x^{5^4+1} \in \mathbb{F}_5[x].$$

Then

$$f^*(x) = x + 4x^5 + 3x^{5^2} + 2x^{5^3} + 2x^{5^4} + 2x^{5^5} + 3x^{5^6} + 4x^{5^7} + x^{5^8}.$$

Using Mathematica [11], we find that

$$(2.9) \quad \deg(f^*, x^{5^m} - x) = \begin{cases} 5^4 & \text{if } m = 13, \\ 5^8 & \text{if } m = 26, \\ 1 & \text{if } 1 \leq m \leq 26, m \neq 13, 26. \end{cases}$$

Thus, the splitting field of f^* over $\mathbb{F}_5$ is $\mathbb{F}_{5^{26}}$. Let m be any positive integer written in the form $m = 2^a 13^b m^*$, where $(m^*, 2 \cdot 13) = 1$. Then by (2.8) and (2.9),

$$(2.10) \quad l_m(f) = \begin{cases} 0 & \text{if } b = 0, \\ 4 & \text{if } b \geq 1, a = 0, \\ 8 & \text{if } b \geq 1, a \geq 1. \end{cases}$$

In the appendix, we give the values of $l_m(f)$, where f is of the form (2.2) with $p^n = 3$, $\alpha \leq 4$ and $p^n = 5$, $\alpha \leq 3$.

3. FROM $S(f, n)$ TO $S(f + bx, n)$

Let $f(x) \in \mathbb{F}_{p^n}[x]$ be as in (2.2).

Lemma 3.1. *Let $b \in \mathbb{F}_{p^n}$. Then*

$$\begin{aligned} & S(f, n) \overline{S(f + bx, n)} \\ &= \begin{cases} p^{n+l_n(f)} e_n(f(x_0)) & \text{if } f^*(x) = b^{p^\alpha} \text{ has a solution } x_0 \in \mathbb{F}_{p^n} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Proof. We have

$$\begin{aligned} & S(f, n) \overline{S(f + bx, n)} \\ &= \sum_{x, y \in \mathbb{F}_{p^n}} e_n(f(x) - f(y) - by) \\ &= \sum_{x, y \in \mathbb{F}_{p^n}} e_n(f(x + y) - f(y) - by) \\ &= \sum_{x, y \in \mathbb{F}_{p^n}} e_n(f(x) + y f^*(x)^{p^{-\alpha}} - by) \quad (\text{by (2.4)}) \\ &= \sum_{x \in \mathbb{F}_{p^n}} e_n(f(x)) \sum_{y \in \mathbb{F}_{p^n}} e_n(y(f^*(x)^{p^{-\alpha}} - y)) \\ &= p^n \sum_{\substack{x \in \mathbb{F}_{p^n} \\ f^*(x) = b^{p^\alpha}}} e_n(f(x)). \end{aligned}$$

If $f^*(x) = b^{p^\alpha}$ has no solution in $\mathbb{F}_{p^n}$, $S(f, n) \overline{S(f + bx, n)} = 0$. If $f^*(x) = b^{p^\alpha}$ has a solution $x_0 \in \mathbb{F}_{p^n}$, then the solution set of $f^*(x) = b^{p^\alpha}$ in $\mathbb{F}_{p^n}$ is the $l_n(f)$ -dimensional affine subspace $x_0 + \{z \in \mathbb{F}_{p^n} : f^*(z) = 0\}$ and $\text{Tr}_n(f(x))$ is a constant on this affine subspace. Therefore,

$$S(f, n) \overline{S(f + bx, n)} = p^{n+l_n(f)} e_n(f(x_0)).$$

□

Corollary 3.2. *We have*

$$S(f + bx, n) = \begin{cases} \overline{e_n(f(x_0))} S(f, n) & \text{if } f^*(x) = b^{p^\alpha} \text{ has a solution } x_0 \in \mathbb{F}_{p^n}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Since $S(f, n) \neq 0$, by Lemma 3.1, we only have to consider the case when $f^*(x) = b^{p^\alpha}$ has a solution $x_0 \in \mathbb{F}_{p^n}$. We have

$$\begin{aligned} S(f + bx, n) &= \frac{p^{n+l_n(f)}}{S(f, n)} \overline{e_n(f(x_0))} \\ &= \frac{p^{n+l_n(f)}}{|S(f, n)|^2} \overline{e_n(f(x_0))} S(f, n) \\ &= \overline{e_n(f(x_0))} S(f, n) \quad (\text{by (2.3)}). \end{aligned}$$

□

4. FROM $S(f, n)$ TO $S(f, q^s n)$, $q \neq 2, p$

In this section, we assume that q is a odd prime with $q \neq p$ and $s \geq 0$.

Lemma 4.1. *In the ring $\mathbb{Z}[\zeta_p]$, we have*

$$(4.1) \quad S(f, q^s n) \equiv \sum_{x \in \mathbb{F}_{p^n}} e_{q^s n}(f(x)) \pmod{q};$$

equivalently,

$$(4.2) \quad t_{q^s n}(f) g_p^{q^s n - l_{q^s n}(f)} p^{l_{q^s n}(f)} \equiv \left(\frac{q}{p}\right)^{s(n-l_n(f))} t_n(f) g_p^{n-l_n(f)} p^{l_n(f)} \pmod{q},$$

where $\left(\frac{q}{p}\right)$ is the Legendre symbol ([7]).

Proof. Write

$$S(f, q^s n) = \sum_{x \in \mathbb{F}_{p^{q^s n}} \setminus \mathbb{F}_{p^n}} e_{q^s n}(f(x)) + \sum_{x \in \mathbb{F}_{p^n}} e_{q^s n}(f(x)).$$

In the above,

$$\sum_{x \in \mathbb{F}_{p^{q^s n}} \setminus \mathbb{F}_{p^n}} e_{q^s n}(f(x)) = \sum_{i=0}^{p-1} |T(i)| \zeta_p^i,$$

where $T(i) = \{x \in \mathbb{F}_{p^{q^s n}} \setminus \mathbb{F}_{p^n} : \text{Tr}_{q^s n}(f(x)) = i\}$. $T(i)$ is a union of $\text{Aut}(\mathbb{F}_{p^{q^s n}}/\mathbb{F}_{p^n})$ -orbits of cardinality > 1 . Since the cardinality of every $\text{Aut}(\mathbb{F}_{p^{q^s n}}/\mathbb{F}_{p^n})$ -orbit is a power of q , we have $|T(i)| \equiv 0 \pmod{q}$ for all $0 \leq i \leq p-1$. Hence,

$$\sum_{x \in \mathbb{F}_{p^{q^s n}} \setminus \mathbb{F}_{p^n}} e_{q^s n}(f(x)) \equiv 0 \pmod{q}.$$

So, (4.1) is proved.

To see the equivalence between (4.1) and (4.2), we compute both sides of (4.1). By (2.3),

$$S(f, q^s n) = t_{q^s n}(f) g_p^{q^s n - l_{q^s n}(f)} p^{l_{q^s n}(f)}.$$

Note that the quadratic forms $q^s \text{Tr}_n(f(x))$ and $\text{Tr}_n(f(x))$ on $\mathbb{F}_{p^n}$ have the same rank but the type of $q^s \text{Tr}_n(f(x))$ equals $\left(\frac{q}{p}\right)^{s(n-l_n(f))}$ times the type of $\text{Tr}_n(f(x))$. Thus,

$$\sum_{x \in \mathbb{F}_{p^n}} e_{q^s n}(f(x)) = \sum_{x \in \mathbb{F}_{p^n}} \zeta_p^{q^s \text{Tr}_n(f(x))} = \left(\frac{q}{p}\right)^{s(n-l_n(f))} t_n(f) g_p^{n-l_n(f)} p^{l_n(f)}.$$

Now, the equivalence between (4.1) and (4.2) is clear. □

Lemma 4.2. *We have*

$$(4.3) \quad p^{l_{q^s n}(f) - l_n(f)} \equiv 1 \pmod{q}$$

and

$$(4.4) \quad \frac{1}{2} [l_{q^s n}(f) - l_n(f)] \in \mathbb{Z}.$$

Proof. Let $X = \{x \in \mathbb{F}_{p^{q^s n}} \setminus \mathbb{F}_{p^n} : f^*(x) = 0\}$. Then X is a union of $\text{Aut}(\mathbb{F}_{p^{q^s n}}/\mathbb{F}_{p^n})$ -orbits of cardinality > 1 . Thus,

$$(4.5) \quad p^{l_{q^s n}(f)} - p^{l_n(f)} = |X| \equiv 0 \pmod{q}$$

and (4.3) is proved.

Using (4.3), we can simplify (4.2) as

$$(4.6) \quad t_{q^s n}(f) g_p^{q^s n - l_{q^s n}(f)} \equiv \left(\frac{q}{p}\right)^{s(n - l_n(f))} t_n(f) g_p^{n - l_n(f)} \pmod{q}.$$

Assume to the contrary of (4.4) that $l_{q^s n}(f) - l_n(f)$ is odd. Then exactly one of $q^s n - l_{q^s n}(f)$ and $n - l_n(f)$ is odd. Also note that $g_p^2 = \pm p$. Thus (4.6) gives

$$(4.7) \quad g_p p^a \equiv b \pmod{q}$$

for some $a, b \in \mathbb{Z}$, $a \geq 0$. Let $\sigma \in \text{Aut}(\mathbb{Q}(g_p)/\mathbb{Q})$ such that $\sigma(g_p) = -g_p$. Since $\mathbb{Q}(\zeta_p)/\mathbb{Q}$ is Galois, we can extend σ to $\tau \in \text{Aut}(\mathbb{Q}(\zeta_p)/\mathbb{Q})$. Apply τ to (4.7). We have

$$-g_p p^a \equiv b \pmod{q}.$$

It follows that $2g_p p^a \equiv 0 \pmod{q}$, which is a contradiction. $\square$

Let $o_q(p)$ denote the multiplicative order of p in $\mathbb{Z}/q\mathbb{Z}$. Then by (4.3), $\frac{1}{o_q(p)}(l_{q^s n}(f) - l_n(f)) \in \mathbb{Z}$.

Theorem 4.3. *We have*

$$(4.8) \quad t_{q^s n}(f) = t_n(f) \left(\frac{q}{p}\right)^{sl_n(f)} (-1)^{\frac{1}{4}(p-1)[l_{q^s n}(f) - l_n(f)] + \frac{1}{o_q(p)}[l_{q^s n}(f) - l_n(f)]}.$$

Proof. By (4.6),

$$(4.9) \quad t_{q^s n}(f) g_p^{n(q^s - 1)} \equiv \left(\frac{q}{p}\right)^{s(n - l_n(f))} t_n(f) g_p^{l_{q^s n}(f) - l_n(f)} \pmod{q}.$$

Since $g_p^2 = (-1)^{\frac{p-1}{2}} p$, we have

$$(4.10) \quad \begin{aligned} g_p^{n(q^s - 1)} &= \left[(-1)^{\frac{p-1}{2}} p\right]^{\frac{1}{2}n(q^s - 1)} = (-1)^{\frac{1}{4}n(p-1)(q^s - 1)} p^{\frac{1}{2}(q-1)n(1+q+\dots+q^{s-1})} \\ &\equiv (-1)^{\frac{1}{4}sn(p-1)(q-1)} \left(\frac{p}{q}\right)^{sn} \pmod{q}. \end{aligned}$$

(Note that $p^{\frac{1}{2}(q-1)} \equiv \left(\frac{p}{q}\right) \pmod{q}$ and that $q^s - 1 \equiv s(q-1) \pmod{4}$.) Also,

$$\begin{aligned} g_p^{l_{q^s n}(f) - l_n(f)} &= \left[(-1)^{\frac{p-1}{2}} p\right]^{\frac{1}{2}(l_{q^s n}(f) - l_n(f))} \\ &= (-1)^{\frac{1}{4}(p-1)(l_{q^s n}(f) - l_n(f))} p^{\frac{1}{2}(l_{q^s n}(f) - l_n(f))}. \end{aligned}$$

Since $\frac{1}{2}(l_{q^s n}(f) - l_n(f)) \in \mathbb{Z}$ and $o_q(p) \mid (l_{q^s n}(f) - l_n(f))$ (Lemma 4.2), we have

$$p^{\frac{1}{2}(l_{q^s n}(f) - l_n(f))} \equiv (-1)^{\frac{1}{o_q(p)}(l_{q^s n}(f) - l_n(f))} \pmod{q}.$$

Thus,

$$(4.11) \quad g_p^{l_{q^n}(f)-l_n(f)} \equiv (-1)^{\frac{1}{4}(p-1)(l_{q^n}(f)-l_n(f))+\frac{1}{o_q(p)}(l_{q^n}(f)-l_n(f))} \pmod{q}.$$

Using (4.10) and (4.11) in (4.9), we have

$$\begin{aligned} t_{q^n}(f) &\equiv t_n(f) \left(\frac{q}{p}\right)^{s(n-l_n(f))} \left(\frac{p}{q}\right)^{sn} \\ &\quad \cdot (-1)^{\frac{1}{4}(p-1)(sn(q-1)+l_{q^n}(f)-l_n(f))+\frac{1}{o_q(p)}(l_{q^n}(f)-l_n(f))} \pmod{q}. \end{aligned}$$

In the above, both sides are ± 1 ; hence the two sides are equal. Using the law of quadratic reciprocity

$$\left(\frac{p}{q}\right)\left(\frac{q}{p}\right) = (-1)^{\frac{1}{4}(p-1)(q-1)},$$

we have (4.8). $\square$

Example 4.4. (Example 2.2 continued) Let

$$f(x) = x^{5^0+1} + 2x^{5^1+1} + 3x^{5^2+1} + 4x^{5^3+1} + x^{5^4+1} \in \mathbb{F}_5[x]$$

be the polynomial considered in Example 2.2. The nullity $l_m(f)$ is given by (2.10) for all $m > 0$. Since

$$S(f, 1) = \sum_{x \in \mathbb{F}_5} e_1(f(x)) = \sum_{x \in \mathbb{F}_5} e_1(x^2) = g_5,$$

we have

$$t_1(f) = 1.$$

Now let m be odd and not divisible by 5. Write $m = 13^b m^*$, where $(m^*, 2 \cdot 5 \cdot 13) = 1$. Note that $o_{13}(5) = 4$. Then it follows from (4.8) that

$$t_m(f) = \begin{cases} 1 & \text{if } b = 0, \\ -1 & \text{if } b \geq 1. \end{cases}$$

In the next section, we will revisit this example and determine $t_m(f)$ for all even m not divisible by 5.

5. FROM $S(f, n)$ TO $S(f, 2^s n)$, $s > 0$

We have

$$S(f, 2^s n) = \sum_{x \in \mathbb{F}_{p^{2^s n}} \setminus \mathbb{F}_{p^{2n}}} e_{2^s n}(f(x)) + \sum_{x \in \mathbb{F}_{p^{2n}}} e_{2^s n}(f(x)).$$

Here,

$$\sum_{x \in \mathbb{F}_{p^{2^s n}} \setminus \mathbb{F}_{p^{2n}}} e_{2^s n}(f(x)) = \sum_{i=0}^{p-1} |T(i)| \zeta_p^i,$$

where

$$T(i) = \{x \in \mathbb{F}_{p^{2^s n}} \setminus \mathbb{F}_{p^{2n}} : \text{Tr}_{2^s n}(f(x)) = i\}.$$

We claim that $T(i)$ is a union of $\text{Aut}(\mathbb{F}_{p^{2^s n}}/\mathbb{F}_{p^n})$ -orbits of cardinality divisible by 4. Note that $\text{Aut}(\mathbb{F}_{p^{2^s n}}/\mathbb{F}_{p^n})$ is cyclic of order 2^s . If $x \in \mathbb{F}_{p^{2^s n}} \setminus \mathbb{F}_{p^{2n}}$, its stabilizer in $\text{Aut}(\mathbb{F}_{p^{2^s n}}/\mathbb{F}_{p^n})$ does not contain $\text{Aut}(\mathbb{F}_{p^{2^s n}}/\mathbb{F}_{p^{2n}})$ hence must be properly contained in $\text{Aut}(\mathbb{F}_{p^{2^s n}}/\mathbb{F}_{p^{2n}})$. So, the $\text{Aut}(\mathbb{F}_{p^{2^s n}}/\mathbb{F}_{p^n})$ -orbit of x has a cardinality divisible by 4; hence the claim is proved.

Therefore, we have

$$(5.1) \quad S(f, 2^s n) \equiv \sum_{x \in \mathbb{F}_{p^{2n}}} e_{2^s n}(f(x)) \pmod{4}.$$

Partition $\mathbb{F}_{p^{2n}}$ as

$$\mathbb{F}_{p^{2n}} = \mathbb{F}_{p^n} \dot{\cup} A \dot{\cup} B,$$

where

$$\begin{aligned} A &= \{x \in \mathbb{F}_{p^{2n}}^* : x^{p^n-1} = -1\}, \\ B &= \{x \in \mathbb{F}_{p^{2n}}^* : x^{p^n-1} \neq \pm 1\}. \end{aligned}$$

Note that B can be further partitioned into four-element subsets of the form $\{\pm x, \pm x^{p^n}\}$. Moreover, $e_{2^s n}(f(x))$ is constant on $\{\pm x, \pm x^{p^n}\}$. Thus,

$$(5.2) \quad \sum_{x \in B} e_{2^s n}(f(x)) \equiv 0 \pmod{4}.$$

Let β be a nonsquare of $\mathbb{F}_{p^n}$ and let $x_0 \in \mathbb{F}_{p^{2n}}$ such that $x_0^2 = \beta$. Then $x_0^{p^n-1} = -1$ and $A = x_0 \mathbb{F}_{p^n}^*$. So,

$$\begin{aligned} \sum_{x \in A} e_{2^s n}(f(x)) &= \sum_{x \in \mathbb{F}_{p^n}^*} e_{2^s n}\left(\sum_{i=1}^k a_i x_0^{p^{\alpha_i}+1} x^{p^{\alpha_i}+1}\right) \\ (5.3) \quad &= \sum_{x \in \mathbb{F}_{p^n}^*} e_{2^s n}\left(\sum_{i=1}^k a_i \beta^{\frac{1}{2}(p^{\alpha_i}+1)} x^{p^{\alpha_i}+1}\right) \\ &= \sum_{x \in \mathbb{F}_{p^n}} e_{2^s n}(\tilde{f}(x)) - 1, \end{aligned}$$

where

$$(5.4) \quad \tilde{f}(x) = \sum_{i=1}^k a_i \beta^{\frac{1}{2}(p^{\alpha_i}+1)} x^{p^{\alpha_i}+1} \in \mathbb{F}_{p^n}[x].$$

Note that when β is fixed, $\tilde{f}(x) = f(\beta x)$. By (5.1) – (5.3), we have

$$\begin{aligned} S(f, 2^s n) &\equiv \sum_{x \in \mathbb{F}_{p^n}} e_{2^s n}(f(x)) + \sum_{x \in A} e_{2^s n}(f(x)) \pmod{4} \\ &= \sum_{x \in \mathbb{F}_{p^n}} e_{2^s n}(f(x)) + \sum_{x \in \mathbb{F}_{p^n}} e_{2^s n}(\tilde{f}(x)) - 1 \\ &= \left(\frac{2}{p}\right)^{s(n-l_n(f))} S(f, n) + \left(\frac{2}{p}\right)^{s(n-l_n(\tilde{f}))} S(\tilde{f}, n) - 1, \end{aligned}$$

i.e.,

$$\begin{aligned} &t_{2^s n}(f) g_p^{2^s n - l_{2^s n}(f)} p^{l_{2^s n}(f)} \\ &\equiv \left(\frac{2}{p}\right)^{s(n-l_n(f))} t_n(f) g_p^{n-l_n(f)} p^{l_n(f)} + \left(\frac{2}{p}\right)^{s(n-l_n(\tilde{f}))} t_n(\tilde{f}) g_p^{n-l_n(\tilde{f})} p^{l_n(\tilde{f})} \pmod{4}. \end{aligned}$$

Since $\left(\frac{2}{p}\right) = (-1)^{\frac{1}{8}(p^2-1)}$ and $p \equiv (-1)^{\frac{1}{2}(p-1)} \pmod{4}$, the above can be written as

$$(5.5) \quad \begin{aligned} & t_{2^s n}(f) (-1)^{\frac{1}{2}(p-1)l_{2^s n}(f)} g_p^{2^s n - l_{2^s n}(f)} \\ & \equiv t_n(f) (-1)^{\frac{1}{8}(p^2-1)s(n-l_n(f)) + \frac{1}{2}(p-1)l_n(f)} g_p^{n-l_n(f)} \\ & \quad + t_n(\tilde{f}) (-1)^{\frac{1}{8}(p^2-1)s(n-l_n(\tilde{f})) + \frac{1}{2}(p-1)l_n(\tilde{f})} g_p^{n-l_n(\tilde{f})} - 1 \pmod{4}. \end{aligned}$$

Theorem 5.1. *Let f be given by (2.2) and $\tilde{f}$ by (5.4) and let $s > 0$. Then $l_n(f) + l_n(\tilde{f}) + l_{2^s n}(f)$ is even. Moreover,*

$$(5.6) \quad t_{2^s n}(f) = \begin{cases} t_n(f)t_n(\tilde{f}) & \text{if } l_n(f) \equiv l_n(\tilde{f}) \pmod{2}, \\ t_n(f)t_n(\tilde{f})(-1)^{\frac{1}{8}(p^2-1)s} & \text{if } l_n(f) \not\equiv l_n(\tilde{f}) \pmod{2}. \end{cases}$$

Proof. We first show that $l_n(f) + l_n(\tilde{f}) + l_{2^s n}(f)$ is even. If, to the contrary, $l_n(f) + l_n(\tilde{f}) + l_{2^s n}(f)$ is odd, then exactly one or three of $2^s n - l_{2^s n}(f)$, $n - l_n(f)$ and $n - l_n(\tilde{f})$ are odd. Note that $g_p^2 = (-1)^{\frac{1}{4}(p-1)^2} p \equiv 1 \pmod{4}$. Thus, (5.5) gives

$$g_p \equiv u \pmod{4} \quad \text{for some } u \in \mathbb{Z}$$

or

$$-1 \equiv v g_p \pmod{4} \quad \text{for some } v \in \mathbb{Z},$$

depending on whether one or three of $2^s n - l_{2^s n}(f)$, $n - l_n(f)$ and $n - l_n(\tilde{f})$ are odd. Let $\tau \in \text{Aut}(\mathbb{Q}(\zeta_p)/\mathbb{Q})$ such that $\tau(g_p) = -g_p$ and apply τ to the above. In the first case, we have

$$\begin{cases} g_p \equiv u \pmod{4}, \\ -g_p \equiv u \pmod{4}. \end{cases}$$

So, $2g_p \equiv 0 \pmod{4}$, which is a contradiction. In the second case, we have

$$\begin{cases} -1 \equiv v g_p \pmod{4}, \\ -1 \equiv -v g_p \pmod{4}. \end{cases}$$

So, $-2 \equiv 0 \pmod{4}$, which is also a contradiction. Thus, we have proved that $l_n(f) + l_n(\tilde{f}) + l_{2^s n}(f)$ is even.

To prove (5.6), we first assume $l_n(f) \equiv l_n(\tilde{f}) \pmod{2}$. Then $l_{2^s n}(f)$ is even. By (5.5), we have

$$(5.7) \quad t_{2^s n}(f) \equiv (t_n(f) + t_n(\tilde{f}))\delta - 1 \pmod{4},$$

where $\delta \in \{\pm 1, \pm g_p\}$. Since $\frac{1}{2}(g_p - 1)$ is integral over $\mathbb{Q}$, $g_p \equiv 1 \pmod{2}$. Thus, $2\delta \equiv 2 \pmod{4}$. If $t_n(f) + t_n(\tilde{f}) = 0$, (5.7) gives $t_{2^s n}(f) = -1$. If $t_n(f) + t_n(\tilde{f}) = \pm 2$, (5.7) gives

$$t_{2^s n}(f) \equiv 2\delta - 1 \equiv 1 \pmod{4},$$

i.e., $t_{2^s n}(f) = 1$. To sum up, we have

$$t_{2^s n}(f) = \begin{cases} -1 & \text{if } t_n(f) + t_n(\tilde{f}) = 0 \\ 1 & \text{if } t_n(f) + t_n(\tilde{f}) = \pm 2 \end{cases} = t_n(f)t_n(\tilde{f}).$$

Now assume $l_n(f) \not\equiv l_n(\tilde{f}) \pmod{2}$. Then $l_{2^s n}(f)$ is odd. Without loss of generality, assume $n - l_n(f)$ is odd and $n - l_n(\tilde{f})$ is even. Then by (5.5),

$$\begin{aligned} & t_{2^s n}(f)(-1)^{\frac{1}{2}(p-1)} g_p \\ & \equiv t_n(f)(-1)^{\frac{1}{8}(p^2-1)s + \frac{1}{2}(p-1)l_n(f)} g_p + t_n(\tilde{f})(-1)^{\frac{1}{2}(p-1)l_n(\tilde{f})} - 1 \pmod{4} \\ & = t_n(f)(-1)^{\frac{1}{8}(p^2-1)s + \frac{1}{2}(p-1)(l_n(\tilde{f})+1)} g_p + t_n(\tilde{f})(-1)^{\frac{1}{2}(p-1)l_n(\tilde{f})} - 1, \end{aligned}$$

i.e.,

$$\begin{aligned} & (-1)^{\frac{1}{2}(p-1)} \left[t_{2^s n}(f) - t_n(f)(-1)^{\frac{1}{8}(p^2-1)s + \frac{1}{2}(p-1)l_n(\tilde{f})} \right] g_p \\ & \equiv t_n(\tilde{f})(-1)^{\frac{1}{2}(p-1)l_n(\tilde{f})} - 1 \pmod{4}. \end{aligned}$$

So,

$$\begin{aligned} t_{2^s n}(f) &= \begin{cases} t_n(f)(-1)^{\frac{1}{8}(p^2-1)s + \frac{1}{2}(p-1)l_n(\tilde{f})} & \text{if } t_n(\tilde{f})(-1)^{\frac{1}{2}(p-1)l_n(\tilde{f})} = 1, \\ -t_n(f)(-1)^{\frac{1}{8}(p^2-1)s + \frac{1}{2}(p-1)l_n(\tilde{f})} & \text{if } t_n(\tilde{f})(-1)^{\frac{1}{2}(p-1)l_n(\tilde{f})} = -1 \end{cases} \\ &= t_n(f)t_n(\tilde{f})(-1)^{\frac{1}{8}(p^2-1)s}. \end{aligned}$$

This completes the proof of the theorem. $\square$

Example 5.2. (Example 4.4 revisited) Recall that

$$f(x) = x^{5^0+1} + 2x^{5^1+1} + 3x^{5^2+1} + 4x^{5^3+1} + x^{5^4+1} \in \mathbb{F}_5[x],$$

and $t_1(f) = 1$, $l_1(f) = 0$. Choose a nonsquare $\beta = 2 \in \mathbb{F}_5$. Then

$$\tilde{f}(x) = 2x^{5^0+1} + x^{5^1+1} + x^{5^2+1} + 2x^{5^3+1} + 2x^{5^4+1}.$$

Since

$$S(\tilde{f}, 1) = \sum_{x \in \mathbb{F}_5} e_1(\tilde{f}(x)) = \sum_{x \in \mathbb{F}_5} e_1(3x^2) = -g_5,$$

we have $t_1(\tilde{f}) = -1$ and $l_1(\tilde{f}) = 0$. By (5.6),

$$t_{2^a}(f) = -1 \quad \text{for } a > 0.$$

Let $m = 2^a 13^b m^*$, where $s > 0$ and $(m^*, 2 \cdot 5 \cdot 13) = 1$. Then by (4.8),

$$t_m(f) = t_{2^a}(f) = -1.$$

Example 5.3. Let

$$f(x) = x^{3^0+1} + 2x^{3^1+1} + 2x^{3^2+1} + 2x^{3^3+1} + x^{3^4+1} \in \mathbb{F}_3[x].$$

Then

$$f^*(x) = x + 2x^3 + 2x^{3^2} + 2x^{3^3} + 2x^{3^4} + 2x^{3^5} + 2x^{3^6} + 2x^{3^7} + x^{3^8}.$$

The splitting field of f^* over $\mathbb{F}_3$ is $\mathbb{F}_{3^{24}}$ and

$$l_{2^a 3^b m^*}(f) = \begin{cases} 0 & \text{if } a = 0, \\ 1 & \text{if } a = 1, b = 0, \\ 2 & \text{if } a = 1, b \geq 1, \\ 3 & \text{if } a = 2, b = 0, \\ 4 & \text{if } a = 2, b \geq 1, \\ 7 & \text{if } a \geq 3, b = 0, \\ 8 & \text{if } a \geq 3, b \geq 1, \end{cases}$$

where $(m^*, 2 \cdot 3) = 1$. (See Table 1 in the appendix.) Since

$$S(f, 1) = \sum_{x \in \mathbb{F}_3} e_1(2x^2) = -g_3,$$

we have $t_1(f) = -1$. By Theorem 4.3,

$$t_{m^*}(f) = t_1(f) = -1 \quad \text{for all } m^* > 0 \text{ with } (m^*, 2 \cdot 3) = 1.$$

Also,

$$\tilde{f}(x) = 2x^{3^0+1} + 2x^{3^1+1} + x^{3^2+1} + 2x^{3^3+1} + 2x^{3^4+1}.$$

Since $S(\tilde{f}, 1) = \sum_{x \in \mathbb{F}_3} e_1(0) = 3$, we have $t_1(\tilde{f}) = 1$ and $l_1(\tilde{f}) = 1$. By Theorem 5.1,

$$t_{2^a}(f) = -(-1)^a = (-1)^{a+1} \quad \text{for all } a > 0.$$

By Theorem 4.3 again,

$$t_{2^a m^*}(f) = (-1)^{a+1} \left(\frac{m^*}{3} \right), \quad a > 0, \quad (m^*, 2 \cdot 3) = 1.$$

Example 5.4. Let

$$f(x) = 5x^{7^0+1} + 6x^{7^1+1} + x^{7^2+1} \in \mathbb{F}_7[x].$$

Using Mathematica, we find that the splitting field of f^* over $\mathbb{F}_7$ is $\mathbb{F}_{7^{56}}$ and

$$l_{2^a 7^b m^*}(f) = \begin{cases} 0 & \text{if } a = 0, \\ 1 & \text{if } 1 \leq a \leq 2, \ b = 0, \\ 2 & \text{if } 1 \leq a \leq 2, \ b \geq 1, \\ 3 & \text{if } a \geq 3, \ b = 0, \\ 4 & \text{if } a \geq 3, \ b \geq 1, \end{cases}$$

where $(m^*, 2 \cdot 7) = 1$. Since

$$S(f, 1) = \sum_{x \in \mathbb{F}_7} e_1(5x^2) = -g_7,$$

we have $t_1(f) = -1$. By Theorem 4.3,

$$t_{m^*}(f) = t_1(f) = -1 \quad \text{for all } m^* > 0 \text{ with } (m^*, 2 \cdot 7) = 1.$$

Choose a nonsquare $\beta = 3 \in \mathbb{F}_7$. Then

$$\tilde{f}(x) = x^{7^0+1} + 3x^{7^1+1} + 3x^{7^2+1}.$$

Since $S(\tilde{f}, 1) = \sum_{x \in \mathbb{F}_7} e_1(0) = 7$, we have $t_1(\tilde{f}) = 1$ and $l_1(\tilde{f}) = 1$. By Theorem 5.1,

$$t_{2^a}(f) = -1 \quad \text{for all } a > 0.$$

By Theorem 4.3 again,

$$t_{2^a m^*}(f) = -\left(\frac{m^*}{7} \right), \quad a > 0, \quad (m^*, 2 \cdot 7) = 1.$$

6. FROM $S(f, n)$ TO $S(f, pn)$

In this section, we consider the relative formula of $S(f, pn)$ in terms of $S(f, n)$. This seems to be a difficult situation; the congruence method in the previous two sections does not work here. We are able to express $S(f, pn)$ in terms of $S(f, n)$ only under a very restrictive condition on f .

Let $b \in \mathbb{F}_{p^n}$ such that $\text{Tr}_n(b) \neq 0$. By the Artin-Schreier theorem (cf. [9, Ch. VI, Theorem 6.4]), $\mathbb{F}_{p^{pn}} = \mathbb{F}_{p^n}(\epsilon)$, where $\epsilon^p = \epsilon + b$, and the roots of $x^p - x - b$ are $\epsilon + j$, $j \in \mathbb{F}_p$. For every integer $t \geq 0$, we have

$$\begin{aligned}
 \text{Tr}_{pn/n}(\epsilon^t) &= \sum_{j \in \mathbb{F}_p} (\epsilon + j)^t = \sum_{j \in \mathbb{F}_p} \sum_{s=0}^t \binom{t}{s} j^s \epsilon^{t-s} \\
 (6.1) \quad &= \sum_{s=0}^t \binom{t}{s} \epsilon^{t-s} \sum_{j \in \mathbb{F}_p} j^s \\
 &= - \sum_{i \geq 0} \binom{t}{i(p-1)} \epsilon^{t-i(p-1)}.
 \end{aligned}$$

Lemma 6.1. *Let u, v be integers such that $0 \leq u, v \leq p-1$. Then*

$$(6.2) \quad \text{Tr}_{pn/n}(\epsilon^{u+v}) = \begin{cases} 0 & \text{if } u+v \neq p-1, 2(p-1), \\ -1 & \text{if } u+v = p-1, 2(p-1). \end{cases}$$

If α is a positive integer, then

$$\begin{aligned}
 \text{Tr}_{pn/n}(\epsilon^{u+vp^\alpha}) &= - \binom{v}{u+v-(p-1)} (b^{p^0} + \dots + b^{p^{\alpha-1}})^{u+v-(p-1)} \\
 (6.3) \quad &+ \begin{cases} 0 & \text{if } u+v \neq 2(p-1), \\ -1 & \text{if } u+v = 2(p-1). \end{cases}
 \end{aligned}$$

Proof. Write $u+v = u' + v'p$, where $0 \leq u', v' \leq p-1$. By (6.1), we have

$$\begin{aligned}
 \text{Tr}_{pn/n}(\epsilon^{u+v}) &= - \binom{u+v}{p-1} \epsilon^{u+v-(p-1)} - \binom{u+v}{2(p-1)} \epsilon^{u+v-2(p-1)} \\
 &= - \binom{u'+v'p}{p-1} \epsilon^{u+v-(p-1)} - \binom{u'+v'p}{(p-2)+1p} \epsilon^{u+v-2(p-1)} \\
 &= - \binom{u'}{p-1} \epsilon^{u+v-(p-1)} - \binom{u'}{p-2} \binom{v'}{1} \epsilon^{u+v-2(p-1)}.
 \end{aligned}$$

Equation (6.2) follows immediately from the above.

Now we prove (6.4). By (6.1),

$$\begin{aligned}
& \text{Tr}_{pn/n}(\epsilon^{u+vp^\alpha}) \\
&= - \sum_{\substack{0 \leq s, t \leq p-1 \\ 0 < s+tp^\alpha \equiv 0 \pmod{(p-1)}}} \binom{u+vp^\alpha}{s+tp^\alpha} \epsilon^{u+vp^\alpha-(s+tp^\alpha)} \\
&= - \sum_{\substack{0 \leq s \leq u \\ 0 \leq t \leq v \\ 0 < s+t \equiv 0 \pmod{(p-1)}}} \binom{u}{s} \binom{v}{t} \epsilon^{u-s+(v-t)p^\alpha} \\
&= - \sum_{\substack{0 \leq s \leq u \\ 0 \leq t \leq v \\ u+v > s+t \equiv u+v \pmod{(p-1)}}} \binom{u}{s} \binom{v}{t} \epsilon^{s+tp^\alpha} \quad (s \mapsto u-s, \ t \mapsto v-t).
\end{aligned}$$

Since $\epsilon^p = \epsilon + b$, we have $\epsilon^{p^\alpha} = \epsilon + b^{p^0} + \dots + b^{p^{\alpha-1}}$. Thus,

$$\begin{aligned}
& \text{Tr}_{pn/n}(\epsilon^{u+vp^\alpha}) \\
&= - \sum_{\substack{0 \leq s \leq u \\ 0 \leq t \leq v \\ u+v > s+t \equiv u+v \pmod{(p-1)}}} \binom{u}{s} \binom{v}{t} \epsilon^s (\epsilon + b^{p^0} + \dots + b^{p^{\alpha-1}})^t \\
&= - \sum_{\substack{0 \leq s \leq u \\ 0 \leq t \leq v \\ u+v > s+t \equiv u+v \pmod{(p-1)}}} \binom{u}{s} \binom{v}{t} \sum_{\tau=0}^t \binom{t}{\tau} (b^{p^0} + \dots + b^{p^{\alpha-1}})^{t-\tau} \epsilon^{s+\tau}.
\end{aligned}$$

In the above sum, $s+\tau \leq s+t \leq u+v-(p-1) \leq p-1$. Since $\text{Tr}_{pn/n}(\epsilon^{u+vp^\alpha}) \in \mathbb{F}_{p^n}$, in the above, we only have to sum the terms with $s+\tau = 0$. Therefore,

$$\begin{aligned}
& \text{Tr}_{pn/n}(\epsilon^{u+vp^\alpha}) \\
&= - \sum_{\substack{0 \leq t \leq v \\ u+v > t \equiv u+v \pmod{(p-1)}}} \binom{v}{t} (b^{p^0} + \dots + b^{p^{\alpha-1}})^t \\
&= - \binom{v}{u+v-(p-1)} (b^{p^0} + \dots + b^{p^{\alpha-1}})^{u+v-(p-1)} + \begin{cases} 0 & \text{if } u+v \neq 2(p-1), \\ -1 & \text{if } u+v = 2(p-1). \end{cases}
\end{aligned}$$

This completes the proof of the lemma. $\square$

Theorem 6.2. *Let f be as in (2.2). Assume $\nu_p(n) < \min\{\nu_p(\alpha_i) : 1 \leq i \leq k\}$. Then,*

$$(6.4) \quad S(f, pn) = p^{\frac{1}{2}(p-3)(n+l_n(f))} |S(f, n)|^2 \overline{S(f, n)}.$$

Moreover,

$$(6.5) \quad l_{pn}(f) = p l_n(f),$$

$$(6.6) \quad t_{pn}(f) = t_n(f).$$

Proof. Let $\nu_p(n) = \nu$ and write $n = p^\nu n'$, $p \nmid n'$. Choose $b \in \mathbb{F}_{p^{p^\nu}}$ such that $\text{Tr}_{p^\nu}(b) \neq 0$. Then $\text{Tr}_n(b) = n' \text{Tr}_{p^\nu}(b) \neq 0$. Since $\nu_p(\alpha_i) > \nu$, we have

$$b^{p^0} + \dots + b^{p^{\alpha_i-1}} = \text{Tr}_{\alpha_i}(b) = 0.$$

(If $\alpha_i = 0$, $b^{p^0} + \cdots + b^{p^{\alpha_i-1}}$ is an empty sum.) By Lemma 6.1, for all $1 \leq i \leq k$ and all $0 \leq u, v \leq p-1$,

$$\mathrm{Tr}_{pn/n}(\epsilon^{u+vp^{\alpha_i}}) = \begin{cases} 0 & \text{if } u+v \neq p-1, \ 2(p-1), \\ -1 & \text{if } u+v = p-1, \ 2(p-1). \end{cases}$$

Let $x = x_0\epsilon^0 + \cdots + x_{p-1}\epsilon^{p-1} \in \mathbb{F}_{p^{pn}}$, where $x_u \in \mathbb{F}_{p^n}$, $0 \leq u \leq p-1$. Then

$$\begin{aligned} & \mathrm{Tr}_{pn}(f(x)) \\ &= \mathrm{Tr}_{pn}\left(\sum_{i=1}^k a_i (x_0\epsilon^0 + \cdots + x_{p-1}\epsilon^{p-1})^{1+p^{\alpha_i}}\right) \\ &= \mathrm{Tr}_{pn}\left(\sum_{i=1}^k a_i \sum_{0 \leq u, v \leq p-1} x_u x_v^{p^{\alpha_i}} \epsilon^{u+vp^{\alpha_i}}\right) \\ &= \mathrm{Tr}_n\left[\sum_{i=1}^k a_i \sum_{0 \leq u, v \leq p-1} x_u x_v^{p^{\alpha_i}} \mathrm{Tr}_{pn/n}(\epsilon^{u+vp^{\alpha_i}})\right] \\ &= -\mathrm{Tr}_n\left[\sum_{i=1}^k a_i \sum_{\substack{0 \leq u, v \leq p-1 \\ u+v=p-1, 2(p-1)}} x_u x_v^{p^{\alpha_i}}\right] \\ &= -\mathrm{Tr}_n\left[\sum_{i=1}^k a_i \left(x_{p-1}^{1+p^{\alpha_i}} + x_0 x_{p-1}^{p^{\alpha_i}} + x_{p-1} x_0^{p^{\alpha_i}} + x_{\frac{1}{2}(p-1)}^{1+p^{\alpha_i}}\right.\right. \\ &\quad \left.\left.+ \sum_{u=1}^{\frac{1}{2}(p-3)} (x_u x_{p-1-u}^{p^{\alpha_i}} + x_{p-1-u} x_u^{p^{\alpha_i}})\right)\right] \\ &= -\mathrm{Tr}_n\left[\sum_{i=1}^k a_i \left((x_0 + x_{p-1})^{1+p^{\alpha_i}} - x_0^{1+p^{\alpha_i}} + x_{\frac{1}{2}(p-1)}^{1+p^{\alpha_i}}\right)\right. \\ &\quad \left.+ \sum_{u=1}^{\frac{1}{2}(p-3)} \sum_{i=1}^k a_i (x_u x_{p-1-u}^{p^{\alpha_i}} + x_{p-1-u} x_u^{p^{\alpha_i}})\right] \\ &= -\mathrm{Tr}_n\left[f(x_0 + x_{p-1}) - f(x_0) + f(x_{\frac{1}{2}(p-1)}) + \sum_{u=1}^{\frac{1}{2}(p-3)} x_u f^*(x_{p-1-u})^{p^{-\alpha_i}}\right]. \end{aligned}$$

Therefore,

$$\begin{aligned}
S(f, pn) &= \sum_{(x_0, \dots, x_{p-1}) \in \mathbb{F}_{p^n}^p} e_n \left[-f(x_0 + x_{p-1}) + f(x_0) - f(x_{\frac{1}{2}(p-1)}) \right. \\
&\quad \left. - \sum_{u=1}^{\frac{1}{2}(p-3)} x_u f^*(x_{p-1-u})^{p^{-\alpha}} \right] \\
&= \overline{S(f, n)}^2 S(f, n) \prod_{u=1}^{\frac{1}{2}(p-3)} \left(\sum_{x_u, x_{p-1-u} \in \mathbb{F}_{p^n}} e_n(-x_u f^*(x_{p-1-u})^{p^{-\alpha}}) \right) \\
&= |S(f, n)|^2 \overline{S(f, n)} \prod_{u=1}^{\frac{1}{2}(p-3)} p^{l_n(f)+n} \\
&= p^{\frac{1}{2}(p-3)(n+l_n(f))} |S(f, n)|^2 \overline{S(f, n)}.
\end{aligned}$$

So, (6.4) is proved.

Equation (6.4) gives

$$t_{pn}(f) g_p^{pn-l_{pn}(f)} p^{l_{pn}(f)} = p^{\frac{1}{2}(p-3)(n+l_n(f))} (p^{\frac{1}{2}(n+l_n(f))})^2 t_n(f) \overline{g_p^{n-l_n(f)}} p^{l_n(f)},$$

i.e.,

$$t_{pn}(f) i^{\frac{1}{4}(p-1)^2(pn-l_{pn}(f))} p^{\frac{1}{2}(pn+l_{pn}(f))} = t_n(f) i^{-\frac{1}{4}(p-1)^2(n-l_n(f))} p^{\frac{1}{2}p(n+l_n(f))}.$$

Thus, $l_{pn}(f) = p l_n(f)$ and

$$\begin{aligned}
t_{pn}(f) &= t_n(f) i^{-\frac{1}{4}(p-1)^2[n-l_n(f)+(pn-l_{pn}(f))]} \\
&= t_n(f) i^{-\frac{1}{4}(p-1)^2(p+1)(n-l_n(f))} \\
&= t_n(f).
\end{aligned}$$

□

Corollary 6.3. *In Theorem 6.2, assume $0 \leq s \leq \min\{\nu_p(\alpha_i) : 1 \leq i \leq k\} - \nu_p(n)$. Then*

$$\begin{aligned}
l_{p^s n}(f) &= p^s l_n(f), \\
t_{p^s n}(f) &= t_n(f).
\end{aligned}$$

Proof. Apply (6.5) and (6.6) repeatedly. □

7. WHEN $\nu_2(\alpha_1) = \dots = \nu_2(\alpha_k)$

Lemma 7.1. *Let $\alpha_1, \dots, \alpha_k \geq 0$ be integers. Then*

$$\gcd(p^{\alpha_1} + 1, \dots, p^{\alpha_k} + 1) > 2 \Leftrightarrow \nu_2(\alpha_1) = \dots = \nu_2(\alpha_k) < \infty.$$

When $\nu_2(\alpha_1) = \dots = \nu_2(\alpha_k) < \infty$,

$$\gcd(p^{\alpha_1} + 1, \dots, p^{\alpha_k} + 1) = p^{\gcd(\alpha_1, \dots, \alpha_k)} + 1.$$

Proof. It suffices to prove the lemma with $k = 2$.

($\Leftarrow$) Since $\frac{\alpha_i}{(\alpha_1, \alpha_2)}$, $i = 1, 2$, are odd, $p^{(\alpha_1, \alpha_2)} + 1 \mid p^{\alpha_i} + 1$ for $i = 1, 2$. Thus, $p^{(\alpha_1, \alpha_2)} + 1 \mid (p^{\alpha_1} + 1, p^{\alpha_2} + 1)$.

On the other hand,

$$\begin{aligned} \frac{1}{2}(p^{\alpha_1} + 1, p^{\alpha_2} + 1) & \mid \frac{1}{2}\left(\frac{1}{2}(p^{2\alpha_1} - 1), \frac{1}{2}(p^{2\alpha_2} - 1)\right) \\ & = \frac{1}{4}(p^{2(\alpha_1, \alpha_2)} - 1) \\ & = \frac{p^{(\alpha_1, \alpha_2)} + 1}{2} \cdot \frac{p^{(\alpha_1, \alpha_2)} - 1}{2}. \end{aligned}$$

Since $(\frac{p^{\alpha_1} + 1}{2}, \frac{p^{(\alpha_1, \alpha_2)} - 1}{2}) = 1$, we must have $\frac{1}{2}(p^{\alpha_1} + 1, p^{\alpha_2} + 1) \mid \frac{1}{2}(p^{(\alpha_1, \alpha_2)} + 1)$, i.e., $(p^{\alpha_1} + 1, p^{\alpha_2} + 1) \mid p^{(\alpha_1, \alpha_2)} + 1$. Hence, $(p^{\alpha_1} + 1, p^{\alpha_2} + 1) = p^{(\alpha_1, \alpha_2)} + 1$.

($\Rightarrow$) Clearly, $\alpha_i > 0$ for all $1 \leq i \leq k$. Assume to the contrary that $\nu_2(\alpha_1) > \nu_2(\alpha_2)$. Write $\alpha_1 = 2^i \alpha'_1$ and $\alpha_2 = 2^j \alpha'_2$, where $i > j$, α'_1 and α'_2 are odd. Then we have

$$\begin{aligned} (p^{\alpha_1} + 1, p^{\alpha_2} + 1) & \mid (p^{2^i \alpha'_1 \alpha'_2} + 1, p^{2\alpha_2} - 1) \\ & \mid (p^{2^i \alpha'_1 \alpha'_2} + 1, p^{2^i \alpha'_1 \alpha'_2} - 1) \\ & = 2, \end{aligned}$$

which is a contradiction. $\square$

Lemma 7.2. *Let $\alpha, \beta \geq 0$ be integers. Then*

$$(p^\alpha + 1, p^\beta - 1) = \begin{cases} p^{(\alpha, \beta)} + 1 & \text{if } \nu_2(\beta) > \nu_2(\alpha), \\ 2 & \text{if } \nu_2(\beta) \leq \nu_2(\alpha). \end{cases}$$

Proof. If any of α and β is 0, the conclusion is obvious. So, assume $\alpha, \beta > 0$.

First assume $\nu_2(\beta) > \nu_2(\alpha)$. Since $\frac{\alpha}{(\alpha, \beta)}$ is odd, $p^{(\alpha, \beta)} + 1 \mid p^\alpha + 1$; since $\frac{\beta}{(\alpha, \beta)}$ is even, $p^{(\alpha, \beta)} + 1 \mid p^\beta - 1$. So, $p^{(\alpha, \beta)} + 1 \mid (p^\alpha + 1, p^\beta - 1)$. Note that

$$\begin{aligned} \frac{1}{2}(p^\alpha + 1, p^\beta - 1) & \mid \frac{1}{2}(p^{2\alpha} - 1, p^\beta - 1) \\ & = \frac{1}{2}(p^{2(\alpha, \beta)} - 1) \\ & = \frac{1}{2}(p^{2(\alpha, \beta)} - 1) \\ & = \frac{1}{2}(p^{(\alpha, \beta)} - 1)(p^{(\alpha, \beta)} + 1). \end{aligned}$$

Since $(\frac{1}{2}(p^\alpha + 1), \frac{1}{2}(p^{(\alpha, \beta)} - 1)) = 1$, we must have

$$(7.1) \quad \frac{1}{2}(p^\alpha + 1, p^\beta - 1) \mid p^{(\alpha, \beta)} + 1.$$

For each $x \in \mathbb{Z}$ and odd integer $k > 0$, we have $\nu_2(1 + x^k) = \nu_2(1 + x)$. By this fact, $\nu_2(p^\alpha + 1) = \nu_2(p^{(\alpha, \beta)} + 1)$ and $\nu_2(p^\beta - 1) \geq \nu_2(p^\alpha + 1)$. So, $\nu_2(p^\alpha + 1, p^\beta - 1) = \nu_2(p^{(\alpha, \beta)} + 1)$. Thus, (7.1) gives $(p^\alpha + 1, p^\beta - 1) \mid p^{(\alpha, \beta)} + 1$. So, we have proved that $(p^\alpha + 1, p^\beta - 1) = p^{(\alpha, \beta)} + 1$.

Now assume $\nu_2(\beta) \leq \nu_2(\alpha)$. We have

$$\begin{aligned} \left(\frac{p^\alpha + 1}{2}, \frac{p^\alpha - 1}{2}, \frac{p^\beta - 1}{2} \right) & \mid \frac{1}{2}(p^{2\alpha} - 1, p^\beta - 1) \\ &= \frac{1}{2}(p^{(2\alpha, \beta)} - 1) \\ &= \frac{1}{2}(p^{(\alpha, \beta)} - 1) \\ &= \left(\frac{p^\alpha - 1}{2}, \frac{p^\beta - 1}{2} \right). \end{aligned}$$

Since $(\frac{p^\alpha + 1}{2}, \frac{p^\alpha - 1}{2}) = 1$, we must have $(\frac{p^\alpha + 1}{2}, \frac{p^\beta - 1}{2}) = 1$, i.e., $(p^\alpha + 1, p^\beta - 1) = 2$. $\square$

Theorem 7.3. *Let f be as in (2.2). Assume that $\nu_2(\alpha_1) = \dots = \nu_2(\alpha_k) = \nu$ and that $\nu_2(n) > \nu$. Then $2^{\nu+1} \mid l_n(f)$ and*

$$t_n(f) = (-1)^{(\frac{1}{4}(p-1)^2 2^\nu + 1) \frac{n - l_n(f)}{2^{\nu+1}}} = \begin{cases} (-1)^{(\frac{1}{4}(p-1)^2 + 1) \frac{n - l_n(f)}{2^{\nu+1}}} & \text{if } \nu = 0, \\ (-1)^{\frac{n - l_n(f)}{2^{\nu+1}}} & \text{if } \nu > 0. \end{cases}$$

Proof. By Lemmas 7.1 and 7.2, we have

$$\begin{aligned} \gcd(p^{\alpha_1} + 1, \dots, p^{\alpha_k} + 1, p^n - 1) &= (p^{\gcd(\alpha_1, \dots, \alpha_k)} + 1, p^n - 1) \\ (7.2) \quad &= p^{\gcd(\alpha_1, \dots, \alpha_k, n)} + 1 \\ &\equiv 0 \pmod{p^{2^\nu} + 1}. \end{aligned}$$

Let $q = p^{2^\nu} + 1$. Then $2^{\nu+1}$ is the multiplicative order of p modulo q , i.e., $o_q(p) = 2^{\nu+1}$. Since $2^{\nu+1} \mid n$, we have $q \mid p^n - 1$.

We first show that $2^{\nu+1} \mid l_n(f)$. In fact, we show that $\{x \in \mathbb{F}_{p^n} : f^*(x) = 0\}$ is a vector space over $\mathbb{F}_{p^{2^\nu+1}}$. Let $x \in \mathbb{F}_{p^n}$ such that $f^*(x) = 0$ and let $y \in \mathbb{F}_{p^{2^\nu+1}}$. We want to show $f^*(yx) = 0$. By (2.5),

$$(7.3) \quad f^*(yx)^{p^{-\alpha}} = \sum_{i=1}^k (a_i y^{p^{\alpha_i}} x^{p^{\alpha_i}} + a_i^{p^{-\alpha_i}} y^{p^{-\alpha_i}} x^{p^{-\alpha_i}}).$$

We claim that $y^{p^{\pm \alpha_i}} = y^{p^\alpha}$ for all $1 \leq i \leq k$. By (7.2), $p^{\alpha_i} \equiv -1 \pmod{q}$; hence $p^{\pm \alpha_i} \equiv -1 \pmod{q}$. Thus, $p^\alpha \equiv -1 \equiv p^{\pm \alpha_i} \pmod{q}$, i.e., $p^{\alpha \mp \alpha_i} \equiv 1 \pmod{q}$ for all $1 \leq i \leq k$. Since $o_q(p) = 2^{\nu+1}$, $\alpha \mp \alpha_i \equiv 0 \pmod{2^{\nu+1}}$. So, $y^{p^{\alpha \mp \alpha_i}} = y$, i.e., $y^{p^\alpha} = y^{p^{\pm \alpha_i}}$. Now by (7.3),

$$f^*(yx)^{p^{-\alpha}} = y^{p^\alpha} \sum_{i=1}^k (a_i x^{p^{\alpha_i}} + a_i^{p^{-\alpha_i}} x^{p^{-\alpha_i}}) = y^{p^\alpha} f^*(x)^{p^{-\alpha}} = 0.$$

Choose $z \in \mathbb{F}_{p^n}^*$ such that $o(z) = q$. Since $p^{\alpha_i} + 1 \equiv 0 \pmod{q}$, $1 \leq i \leq k$, we have

$$f(yx) = f(x) \quad \text{for all } y \in \langle z \rangle.$$

Thus,

$$\begin{aligned}
 t_n(f)g_p^{n-l_n(f)}p^{l_n(f)} &= S(f, n) \\
 &= 1 + \sum_{x \in \mathbb{F}_{p^n}^*} e_n(f(x)) \\
 (7.4) \quad &= 1 + q \sum_{x \in \mathbb{F}_{p^n}^* / \langle z \rangle} e_n(f(x)) \\
 &\equiv 1 \pmod{q}.
 \end{aligned}$$

In the above, $p^{l_n(f)} \equiv 1 \pmod{q}$ since $o_q(p) = 2^{\nu+1} \mid l_n(f)$. Also,

$$g_p^{2^{\nu+1}} = \left[i^{\frac{1}{4}(p-1)^2} p^{\frac{1}{2}} \right]^{2^{\nu+1}} = (-1)^{\frac{1}{4}(p-1)^2 2^{\nu}} p^{2^{\nu}} \equiv (-1)^{\frac{1}{4}(p-1)^2 2^{\nu} + 1} \pmod{q};$$

hence,

$$g_p^{n-l_n(f)} = g_p^{2^{\nu+1} \frac{n-l_n(f)}{2^{\nu+1}}} \equiv (-1)^{(\frac{1}{4}(p-1)^2 2^{\nu} + 1) \frac{n-l_n(f)}{2^{\nu+1}}} \pmod{q}.$$

Now (7.4) gives

$$t_n(f) = (-1)^{(\frac{1}{4}(p-1)^2 2^{\nu} + 1) \frac{n-l_n(f)}{2^{\nu+1}}}.$$

□

Corollary 7.4. Assume $p \equiv -1 \pmod{4}$, $\alpha_1, \dots, \alpha_k$ are all odd and n is even. Then $t_n(f) = 1$.

Proof. This immediate from Theorem 7.3. □

Example 7.5. Let $f(x) = 3x^{5^1+1} + x^{5^3+1} \in \mathbb{F}_5[x]$. Then $f^*(x) = x + 3x^{5^2} + 3x^{5^4} + x^{5^6}$. The splitting field of f^* over $\mathbb{F}_5$ is $\mathbb{F}_{5^{20}}$ and

$$l_{2^a 5^b m^*}(f) = \begin{cases} 0 & \text{if } a \leq 1, \\ 2 & \text{if } a \geq 2, b = 0, \\ 6 & \text{if } a \geq 2, b = 1, \end{cases}$$

where $(m^*, 2 \cdot 5) = 1$. By Theorem 7.3, $t_n(f) = -1$ for all even n .

Theorem 7.3 provides a quick proof for the explicit evaluation of the sum $S(ax^{p^\alpha+1}, n)$.

Corollary 7.6. Let $a \in \mathbb{F}_{p^n}^*$ and let $\alpha \geq 0$.

(i) If $\nu_2(n) \leq \nu_2(\alpha)$,

$$S(ax^{p^\alpha+1}, n) = \eta(a)(-1)^{n-1} i^{\frac{1}{4}(p-1)^2 n} p^{\frac{1}{2}n}.$$

(ii) If $\nu_2(n) = \nu_2(\alpha) + 1$,

$$S(ax^{p^\alpha+1}, n) = \begin{cases} p^{\frac{1}{2}[n+(2\alpha, n)]} & \text{if } a^{\frac{(p^\alpha-1)(p^n-1)}{p^{(2\alpha, n)}-1}} = -1, \\ -p^{\frac{1}{2}n} & \text{otherwise.} \end{cases}$$

(iii) If $\nu_2(n) > \nu_2(\alpha) + 1$,

$$S(ax^{p^\alpha+1}, n) = \begin{cases} -p^{\frac{1}{2}[n+(2\alpha, n)]} & \text{if } a^{\frac{(p^\alpha-1)(p^n-1)}{p^{(2\alpha, n)}-1}} = 1, \\ p^{\frac{1}{2}n} & \text{otherwise.} \end{cases}$$

Proof. We first show that

$$(7.5) \quad l_n(f) = \begin{cases} (2\alpha, n) & \text{if } a^{\frac{(p^\alpha-1)(p^n-1)}{p^{(2\alpha,n)}-1}} = (-1)^{\frac{p^n-1}{p^{(2\alpha,n)}-1}}, \\ 0 & \text{otherwise.} \end{cases}$$

Note that $f^*(x) = a^{p^\alpha} x^{p^{2\alpha}} + ax = a^{p^\alpha} x(x^{p^{2\alpha}-1} + a^{1-p^\alpha})$. Thus, if $f^*(x) = 0$ has a solution in $\mathbb{F}_{p^n}^*$, the number of solutions is $(p^{2\alpha} - 1, p^n - 1) = p^{(2\alpha,n)} - 1$. So,

$$l_n(f) = \begin{cases} (2\alpha, n) & \text{if } f^*(x) = 0 \text{ has a solution in } \mathbb{F}_{p^n}^*, \\ 0 & \text{otherwise.} \end{cases}$$

Observe that $f^*(x) = 0$ has a solution in $\mathbb{F}_{p^n}^*$ if and only if $-a^{p^\alpha-1} = x^{p^{2\alpha}-1}$ for some $x \in \mathbb{F}_{p^n}^*$; the latter holds if and only if

$$\left(-a^{p^\alpha-1}\right)^{\frac{p^n-1}{(p^{2\alpha}-1, p^n-1)}} = 1,$$

i.e.,

$$a^{\frac{(p^\alpha-1)(p^n-1)}{p^{(2\alpha,n)}-1}} = (-1)^{\frac{p^n-1}{p^{(2\alpha,n)}-1}}.$$

So, (7.5) is proved.

(i) Since $\nu_2(n) \leq \nu_2(\alpha)$, by Lemma 7.2, $(p^\alpha+1, p^n-1) = 2$. Therefore, $x \mapsto x^{p^\alpha+1}$ is a 2-to-1 map from $\mathbb{F}_{p^n}^*$ to $(\mathbb{F}_{p^n}^*)^2$. Hence,

$$\begin{aligned} & S(ax^{p^\alpha+1}, n) \\ &= 1 + \sum_{x \in \mathbb{F}_{p^n}^*} e_n(ax^{p^\alpha+1}) = 1 + 2 \sum_{x \in (\mathbb{F}_{p^n}^*)^2} e_n(ax) \\ &= 1 + \sum_{x \in \mathbb{F}_{p^n}^*} e_n(ax^2) = \sum_{x \in \mathbb{F}_{p^n}^*} e_n(ax^2) = \eta(a) \sum_{x \in \mathbb{F}_{p^n}^*} e_n(x^2) \\ &= \eta(a)(-1)^{n-1} g_p^n \quad (\text{by the Davenport-Hasse theorem [4], [10, §5.2]}) \\ &= \eta(a)(-1)^{n-1} i^{\frac{1}{4}(p-1)^2 n} p^{\frac{1}{2}n}. \end{aligned}$$

(ii) Since $\nu_2(n) = \nu_2((2\alpha, n))$, $\frac{p^n-1}{p^{(2\alpha,n)}-1}$ is odd. By (7.5),

$$l_n(f) = \begin{cases} (2\alpha, n) & \text{if } a^{\frac{(p^\alpha-1)(p^n-1)}{p^{(2\alpha,n)}-1}} = -1, \\ 0 & \text{otherwise.} \end{cases}$$

By Theorem 7.3,

$$t_n(f) = (-1)^{(\frac{1}{4}(p-1)^2 \alpha + 1) \frac{n-l_n(f)}{2\nu_2(\alpha)+1}}.$$

Thus,

$$\begin{aligned} S(ax^{p^\alpha+1}, n) &= t_n(f) g_p^{n-l_n(f)} p^{l_n(f)} \\ &= \begin{cases} p^{\frac{1}{2}[n+(2\alpha,n)]} & \text{if } a^{\frac{(p^\alpha-1)(p^n-1)}{p^{(2\alpha,n)}-1}} = -1, \\ -p^{\frac{1}{2}n} & \text{otherwise.} \end{cases} \end{aligned}$$

(In the above, we used the fact that $\nu_2(n - (2\alpha, n)) > \nu_2(n)$ and $\nu_2(2\alpha + n) > \nu_2(n)$.)

(iii) In this case, $\frac{p^n-1}{p^{(2\alpha,n)}-1}$ is even. By (7.5),

$$l_n(f) = \begin{cases} (2\alpha, n) & \text{if } a^{\frac{(p^\alpha-1)(p^n-1)}{p^{(2\alpha,n)}-1}} = 1, \\ 0 & \text{otherwise.} \end{cases}$$

By Theorem 7.3,

$$t_n(f) = (-1)^{(\frac{1}{4}(p-1)^2\alpha+1)\frac{n-l_n(f)}{2\nu_2(\alpha)+1}}.$$

The conclusion follows the same way as in (ii). (Note that in this case, $\nu_2(n - (2\alpha, n)) = \nu_2(\alpha) + 1$.) $\square$

APPENDIX. THE NULLITY $l_m(f)$ WITH $p^n = 3$, $\alpha \leq 4$ AND $p^n = 5$, $\alpha \leq 3$

Let $f(x) = \sum_{i=0}^k a_i x^{p^i+1} \in \mathbb{F}_{p^n}[x]$ with $a_k \in \mathbb{F}_p^*$ and let $0 < m \equiv 0 \pmod{n}$. Since $\text{Tr}_m(a_k^{-1}f) = a_k^{-1}\text{Tr}_m(f)$, we have $l_m(f) = l_m(a_k^{-1}f)$, where $a_k^{-1}f$ is monic. So, when computing $l_m(f)$ with $a_k \in \mathbb{F}_p^*$, we may assume $a_k = 1$.

This appendix contains two tables. Table 1 gives the values of $l_m(f)$ with $p^n = 3$ and $\alpha \leq 4$; table 2 gives the values of $l_m(f)$ with $p^n = 5$ and $\alpha \leq 3$. In both tables, the left column contains the coefficients $a_0, \dots, a_k$ of $f(x) = \sum_{i=0}^k a_i x^{p^i+1}$ with $a_k = 1$. The middle column is s , where $\mathbb{F}_{p^s}$ is the splitting field of $f^*(x)$. The right column lists all pairs $(m, l_m(f))$ such that $m \mid s$. The values of $l_m(f)$ for arbitrary m follows from (2.8), cf. Example 2.2.

REFERENCES

- [1] L. Baumert and R. McEliece, *Weights of irreducible cyclic codes*, Inform. Control **20** (1972), 158 – 175.
- [2] L. Carlitz, *Explicit evaluation of certain exponential sums*, Math. Scand. **44** (1979), 5 – 16.
- [3] L. Carlitz, *Evaluation of some exponential sums over a finite field*, Math. Nachr. **96** (1980), 319 – 339.
- [4] H. Davenport and H. Hasse, *Die Nullstellen der Kongruenzzetafunktionen in gewissen zyklischen Fällen*, J. Reine Angew. Math. **172** (1935), 151 – 182.
- [5] T. Hellese, *Some results about the cross-correlation function between two maximal linear sequences*, Discrete Math. **16** (1976), 209 – 232.
- [6] X. Hou, *Explicit evaluation of certain exponential sums of binary quadratic functions*, preprint.
- [7] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer, New York, 1982.
- [8] N. Jacobson, *Basic Algebra*, W. H. Freeman and Co., New York, 1985.
- [9] S. Lang, *Algebra*, Addison-Wesley, Reading, MA, 1993.
- [10] R. Lidl and H. Niederreiter, *Finite fields*, Cambridge University Press, Cambridge, 1997.
- [11] <http://www.wolfram.com/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH FLORIDA, TAMPA, FL 33620
E-mail address: sdraper95@yahoo.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH FLORIDA, TAMPA, FL 33620
E-mail address: xhou@math.usf.edu

TABLE 1. Values of $l_m(f)$ with $p^n = 3$, $\alpha \leq 4$

$a_0, \dots, a_k$	s	$(m, l_m(f)), m \mid s$
1	1	(1,0)
0 1	4	(1,0) (2,0) (4,2)
1 1	6	(1,0) (2,1) (3,0) (6,2)
2 1	3	(1,1) (3,2)
0 0 1	8	(1,0) (2,0) (4,0) (8,4)
1 0 1	12	(1,0) (2,0) (3,0) (4,2) (6,0) (12,4)
2 0 1	6	(1,1) (2,2) (3,2) (6,4)
0 1 1	18	(1,0) (2,1) (3,0) (6,3) (9,0) (18,4)
1 1 1	12	(1,1) (2,1) (3,2) (4,3) (6,2) (12,4)
2 1 1	5	(1,0) (5,4)
0 2 1	9	(1,1) (3,3) (9,4)
1 2 1	12	(1,0) (2,1) (3,0) (4,3) (6,2) (12,4)
2 2 1	10	(1,0) (2,0) (5,0) (10,4)
0 0 0 1	12	(1,0) (2,0) (3,0) (4,2) (6,0) (12,6)
1 0 0 1	18	(1,0) (2,1) (3,0) (6,3) (9,0) (18,6)
2 0 0 1	9	(1,1) (3,3) (9,6)
0 1 0 1	8	(1,0) (2,0) (4,2) (8,6)
1 1 0 1	30	(1,1) (2,1) (3,2) (5,1) (6,2) (10,5) (15,2) (30,6)
2 1 0 1	30	(1,0) (2,1) (3,0) (5,4) (6,2) (10,5) (15,4) (30,6)
0 2 0 1	12	(1,1) (2,2) (3,2) (4,4) (6,4) (12,6)
1 2 0 1	13	(1,0) (13,6)
2 2 0 1	26	(1,0) (2,0) (13,0) (26,6)
0 0 1 1	30	(1,0) (2,1) (3,0) (5,0) (6,2) (10,5) (15,0) (30,6)
1 0 1 1	12	(1,1) (2,1) (3,2) (4,3) (6,2) (12,6)
2 0 1 1	28	(1,0) (2,0) (4,0) (7,0) (14,0) (28,6)
0 1 1 1	24	(1,1) (2,1) (3,2) (4,1) (6,2) (8,5) (12,2) (24,6)
1 1 1 1	36	(1,0) (2,1) (3,0) (4,3) (6,3) (9,0) (12,5) (18,4) (36,6)
2 1 1 1	7	(1,0) (7,6)
0 2 1 1	13	(1,0) (13,6)
1 2 1 1	20	(1,0) (2,0) (4,2) (5,4) (10,4) (20,6)
2 2 1 1	18	(1,1) (2,2) (3,3) (6,5) (9,4) (18,6)
0 0 2 1	15	(1,1) (3,2) (5,5) (15,6)
1 0 2 1	28	(1,0) (2,0) (4,0) (7,0) (14,0) (28,6)
2 0 2 1	12	(1,0) (2,1) (3,0) (4,3) (6,2) (12,6)
0 1 2 1	24	(1,0) (2,1) (3,0) (4,1) (6,2) (8,5) (12,2) (24,6)
1 1 2 1	14	(1,0) (2,0) (7,0) (14,6)
2 1 2 1	36	(1,1) (2,1) (3,3) (4,3) (6,3) (9,4) (12,5) (18,4) (36,6)
0 2 2 1	26	(1,0) (2,0) (13,0) (26,6)
1 2 2 1	18	(1,1) (2,2) (3,2) (6,5) (9,2) (18,6)
2 2 2 1	20	(1,0) (2,0) (4,2) (5,0) (10,4) (20,6)

TABLE 1. Continued

$a_0, \dots, a_k$	s	$(m, l_m(f)), m \mid s$
0 0 0 0 1	16	(1,0) (2,0) (4,0) (8,0) (16,8)
1 0 0 0 1	24	(1,0) (2,0) (3,0) (4,0) (6,0) (8,4) (12,0) (24,8)
2 0 0 0 1	12	(1,1) (2,2) (3,2) (4,4) (6,4) (12,8)
0 1 0 0 1	90	(1,0) (2,1) (3,0) (5,0) (6,3) (9,0) (10,5) (15,0) (18,4) (30,7) (45, 0) (90,8)
1 1 0 0 1	84	(1,1) (2,1) (3,2) (4,1) (6,2) (7,1) (12,2) (14,1) (21,2) (28,7) (42, 2) (84,8)
2 1 0 0 1	52	(1,0) (2,0) (4,2) (13,0) (26,6) (52,8)
0 2 0 0 1	45	(1,1) (3,3) (5,5) (9,4) (15,7) (45,8)
1 2 0 0 1	84	(1,0) (2,1) (3,0) (4,1) (6,2) (7,0) (12,2) (14,1) (21,0) (28,7) (42, 2) (84,8)
2 2 0 0 1	52	(1,0) (2,0) (4,2) (13,6) (26,6) (52,8)
0 0 1 0 1	36	(1,0) (2,0) (3,0) (4,2) (6,0) (9,0) (12,6) (18,0) (36,8)
1 0 1 0 1	24	(1,1) (2,2) (3,2) (4,2) (6,4) (8,6) (12,4) (24,8)
2 0 1 0 1	10	(1,0) (2,0) (5,4) (10,8)
0 1 1 0 1	36	(1,1) (2,1) (3,3) (4,3) (6,3) (9,6) (12,5) (18,6) (36,8)
1 1 1 0 1	41	(1,0) (41,8)
2 1 1 0 1	42	(1,0) (2,1) (3,0) (6,2) (7,6) (14,7) (21,6) (42,8)
0 2 1 0 1	36	(1,0) (2,1) (3,0) (4,3) (6,3) (9,0) (12,5) (18,6) (36,8)
1 2 1 0 1	82	(1,0) (2,0) (41,0) (82,8)
2 2 1 0 1	42	(1,1) (2,1) (3,2) (6,2) (7,1) (14,7) (21,2) (42,8)
0 0 2 0 1	18	(1,1) (2,2) (3,3) (6,6) (9,4) (18,8)
1 0 2 0 1	24	(1,0) (2,0) (3,0) (4,2) (6,0) (8,6) (12,4) (24,8)
2 0 2 0 1	20	(1,0) (2,0) (4,0) (5,0) (10,0) (20,8)
0 1 2 0 1	41	(1,0) (41,8)
1 1 2 0 1	60	(1,0) (2,1) (3,0) (4,3) (5,4) (6,2) (10,5) (12,4) (15,4) (20,7) (30, 6) (60,8)
2 1 2 0 1	78	(1,1) (2,1) (3,2) (6,2) (13,1) (26,7) (39,2) (78,8)
0 2 2 0 1	82	(1,0) (2,0) (41,0) (82,8)
1 2 2 0 1	60	(1,1) (2,1) (3,2) (4,3) (5,1) (6,2) (10,5) (12,4) (15,2) (20,7) (30, 6) (60,8)
2 2 2 0 1	78	(1,0) (2,1) (3,0) (6,2) (13,6) (26,7) (39,6) (78,8)
0 0 0 1 1	42	(1,0) (2,1) (3,0) (6,2) (7,0) (14,7) (21,0) (42,8)
1 0 0 1 1	78	(1,1) (2,1) (3,2) (6,2) (13,1) (26,7) (39,2) (78,8)
2 0 0 1 1	60	(1,0) (2,0) (3,0) (4,2) (5,4) (6,0) (10,4) (12,4) (15,4) (20,6) (30, 4) (60,8)
0 1 0 1 1	39	(1,1) (3,2) (13,7) (39,8)
1 1 0 1 1	72	(1,0) (2,1) (3,0) (4,1) (6,3) (8,5) (9,0) (12,3) (18,4) (24,7) (36, 4) (72,8)
2 1 0 1 1	52	(1,0) (2,0) (4,2) (13,6) (26,6) (52,8)
0 2 0 1 1	80	(1,0) (2,0) (4,0) (5,0) (8,0) (10,0) (16,0) (20,0) (40,0) (80,8)
1 2 0 1 1	30	(1,0) (2,0) (3,0) (5,0) (6,0) (10,4) (15,0) (30,8)
2 2 0 1 1	36	(1,1) (2,2) (3,3) (4,4) (6,5) (9,4) (12,7) (18,6) (36,8)
0 0 1 1 1	12	(1,1) (2,1) (3,2) (4,3) (6,2) (12,8)
1 0 1 1 1	41	(1,0) (41,8)
2 0 1 1 1	78	(1,0) (2,1) (3,0) (6,2) (13,0) (26,7) (39,0) (78,8)
0 1 1 1 1	60	(1,0) (2,1) (3,0) (4,3) (5,0) (6,2) (10,5) (12,4) (15,0) (20,7) (30, 6) (60,8)
1 1 1 1 1	40	(1,0) (2,0) (4,0) (5,4) (8,4) (10,4) (20,4) (40,8)

TABLE 1. Continued

$a_0, \dots, a_k$	s	$(m, l_m(f)), m \mid s$
2 1 1 1 1	9	(1,1) (3,3) (9,8)
0 2 1 1 1	28	(1,0) (2,0) (4,2) (7,0) (14,0) (28,8)
1 2 1 1 1	18	(1,1) (2,2) (3,2) (6,5) (9,2) (18,8)
2 2 1 1 1	80	(1,0) (2,0) (4,0) (5,0) (8,0) (10,0) (16,0) (20,0) (40,0) (80,8)
0 0 2 1 1	41	(1,0) (41,8)
1 0 2 1 1	36	(1,0) (2,1) (3,0) (4,3) (6,3) (9,0) (12,7) (18,4) (36,8)
2 0 2 1 1	90	(1,1) (2,1) (3,3) (5,1) (6,3) (9,4) (10,5) (15,3) (18,4) (30,7) (45, 4) (90,8)
0 1 2 1 1	82	(1,0) (2,0) (41,0) (82,8)
1 1 2 1 1	24	(1,1) (2,1) (3,2) (4,3) (6,2) (8,7) (12,4) (24,8)
2 1 2 1 1	84	(1,0) (2,1) (3,0) (4,1) (6,2) (7,0) (12,2) (14,1) (21,0) (28,7) (42, 2) (84,8)
0 2 2 1 1	30	(1,1) (2,2) (3,2) (5,5) (6,4) (10,6) (15,6) (30,8)
1 2 2 1 1	28	(1,0) (2,0) (4,2) (7,6) (14,6) (28,8)
2 2 2 1 1	40	(1,0) (2,0) (4,0) (5,0) (8,0) (10,0) (20,0) (40,8)
0 0 0 2 1	21	(1,1) (3,2) (7,7) (21,8)
1 0 0 2 1	78	(1,0) (2,1) (3,0) (6,2) (13,6) (26,7) (39,6) (78,8)
2 0 0 2 1	60	(1,0) (2,0) (3,0) (4,2) (5,0) (6,0) (10,4) (12,4) (15,0) (20,6) (30, 4) (60,8)
0 1 0 2 1	80	(1,0) (2,0) (4,0) (5,0) (8,0) (10,0) (16,0) (20,0) (40,0) (80,8)
1 1 0 2 1	15	(1,0) (3,0) (5,4) (15,8)
2 1 0 2 1	36	(1,1) (2,2) (3,2) (4,4) (6,5) (9,2) (12,7) (18,6) (36,8)
0 2 0 2 1	78	(1,0) (2,1) (3,0) (6,2) (13,0) (26,7) (39,0) (78,8)
1 2 0 2 1	72	(1,1) (2,1) (3,3) (4,1) (6,3) (8,5) (9,4) (12,3) (18,4) (24,7) (36, 4) (72,8)
2 2 0 2 1	52	(1,0) (2,0) (4,2) (13,0) (26,6) (52,8)
0 0 1 2 1	12	(1,0) (2,1) (3,0) (4,3) (6,2) (12,8)
1 0 1 2 1	82	(1,0) (2,0) (41,0) (82,8)
2 0 1 2 1	39	(1,1) (3,2) (13,7) (39,8)
0 1 1 2 1	28	(1,0) (2,0) (4,2) (7,0) (14,0) (28,8)
1 1 1 2 1	18	(1,1) (2,2) (3,3) (6,5) (9,6) (18,8)
2 1 1 2 1	80	(1,0) (2,0) (4,0) (5,0) (8,0) (10,0) (16,0) (20,0) (40,0) (80,8)
0 2 1 2 1	60	(1,1) (2,1) (3,2) (4,3) (5,5) (6,2) (10,5) (12,4) (15,6) (20,7) (30, 6) (60,8)
1 2 1 2 1	40	(1,0) (2,0) (4,0) (5,0) (8,4) (10,4) (20,4) (40,8)
2 2 1 2 1	18	(1,0) (2,1) (3,0) (6,3) (9,0) (18,8)
0 0 2 2 1	82	(1,0) (2,0) (41,0) (82,8)
1 0 2 2 1	36	(1,1) (2,1) (3,3) (4,3) (6,3) (9,4) (12,7) (18,4) (36,8)
2 0 2 2 1	90	(1,0) (2,1) (3,0) (5,4) (6,3) (9,0) (10,5) (15,4) (18,4) (30,7) (45, 4) (90,8)
0 1 2 2 1	30	(1,1) (2,2) (3,2) (5,1) (6,4) (10,6) (15,2) (30,8)
1 1 2 2 1	28	(1,0) (2,0) (4,2) (7,0) (14,6) (28,8)
2 1 2 2 1	40	(1,0) (2,0) (4,0) (5,0) (8,0) (10,0) (20,0) (40,8)
0 2 2 2 1	41	(1,0) (41,8)
1 2 2 2 1	24	(1,0) (2,1) (3,0) (4,3) (6,2) (8,7) (12,4) (24,8)
2 2 2 2 1	84	(1,1) (2,1) (3,2) (4,1) (6,2) (7,1) (12,2) (14,1) (21,2) (28,7) (42, 2) (84,8)

TABLE 2. Values of $l_m(f)$ with $p^n = 5$, $\alpha \leq 3$

$a_0, \dots, a_k$	s	$(m, l_m(f)), m \mid s$
1	1	(1,0)
0 1	4	(1,0) (2,0) (4,2)
1 1	10	(1,0) (2,1) (5,0) (10,2)
2 1	6	(1,0) (2,0) (3,0) (6,2)
3 1	3	(1,0) (3,2)
4 1	5	(1,1) (5,2)
0 0 1	8	(1,0) (2,0) (4,0) (8,4)
1 0 1	20	(1,0) (2,0) (4,2) (5,0) (10,0) (20,4)
2 0 1	12	(1,0) (2,0) (3,0) (4,0) (6,0) (12,4)
3 0 1	6	(1,0) (2,0) (3,2) (6,4)
4 0 1	10	(1,1) (2,2) (5,2) (10,4)
0 1 1	30	(1,0) (2,1) (3,0) (5,0) (6,3) (10,2) (15,0) (30,4)
1 1 1	12	(1,0) (2,0) (3,2) (4,2) (6,2) (12,4)
2 1 1	13	(1,0) (13,4)
3 1 1	5	(1,1) (5,4)
4 1 1	24	(1,0) (2,0) (3,0) (4,0) (6,0) (8,0) (12,0) (24,4)
0 2 1	26	(1,0) (2,0) (13,0) (26,4)
1 2 1	20	(1,0) (2,1) (4,3) (5,0) (10,2) (20,4)
2 2 1	30	(1,1) (2,1) (3,1) (5,2) (6,3) (10,2) (15,2) (30,4)
3 2 1	13	(1,0) (13,4)
4 2 1	15	(1,0) (3,2) (5,0) (15,4)
0 3 1	13	(1,0) (13,4)
1 3 1	20	(1,1) (2,1) (4,3) (5,2) (10,2) (20,4)
2 3 1	30	(1,0) (2,1) (3,2) (5,0) (6,3) (10,2) (15,2) (30,4)
3 3 1	26	(1,0) (2,0) (13,0) (26,4)
4 3 1	30	(1,0) (2,0) (3,0) (5,0) (6,2) (10,0) (15,0) (30,4)
0 4 1	15	(1,1) (3,3) (5,2) (15,4)
1 4 1	12	(1,0) (2,0) (3,0) (4,2) (6,2) (12,4)
2 4 1	26	(1,0) (2,0) (13,0) (26,4)
3 4 1	10	(1,0) (2,1) (5,0) (10,4)
4 4 1	24	(1,0) (2,0) (3,0) (4,0) (6,0) (8,0) (12,0) (24,4)
0 0 0 1	12	(1,0) (2,0) (3,0) (4,2) (6,0) (12,6)
1 0 0 1	30	(1,0) (2,1) (3,0) (5,0) (6,3) (10,2) (15,0) (30,6)
2 0 0 1	18	(1,0) (2,0) (3,0) (6,0) (9,0) (18,6)
3 0 0 1	9	(1,0) (3,0) (9,6)
4 0 0 1	15	(1,1) (3,3) (5,2) (15,6)
0 1 0 1	8	(1,0) (2,0) (4,2) (8,6)
1 1 0 1	126	(1,0) (2,0) (3,0) (6,0) (7,0) (9,0) (14,0) (18,0) (21,0) (42,0) (63, 0) (126,6)
2 1 0 1	30	(1,0) (2,1) (3,2) (5,0) (6,3) (10,4) (15,2) (30,6)
3 1 0 1	30	(1,1) (2,1) (3,1) (5,4) (6,3) (10,4) (15,4) (30,6)
4 1 0 1	63	(1,0) (3,0) (7,0) (9,0) (21,0) (63,6)
0 2 0 1	12	(1,0) (2,0) (3,2) (4,2) (6,4) (12,6)

TABLE 2. Continued

$a_0, \dots, a_k$	s	$(m, l_m(f)), m \mid s$
1 2 0 1	21	(1,0) (3,0) (7,0) (21,6)
2 2 0 1	130	(1,1) (2,1) (5,2) (10,2) (13,1) (26,5) (65,2) (130,6)
3 2 0 1	130	(1,0) (2,1) (5,0) (10,2) (13,4) (26,5) (65,4) (130,6)
4 2 0 1	42	(1,0) (2,0) (3,0) (6,0) (7,0) (14,0) (21,0) (42,6)
0 3 0 1	20	(1,0) (2,0) (4,2) (5,0) (10,0) (20,6)
1 3 0 1	65	(1,1) (5,2) (13,5) (65,6)
2 3 0 1	24	(1,0) (2,0) (3,0) (4,0) (6,2) (8,0) (12,2) (24,6)
3 3 0 1	24	(1,0) (2,0) (3,2) (4,0) (6,2) (8,0) (12,2) (24,6)
4 3 0 1	130	(1,0) (2,1) (5,0) (10,2) (13,0) (26,5) (65,0) (130,6)
0 4 0 1	20	(1,1) (2,2) (4,4) (5,2) (10,4) (20,6)
1 4 0 1	78	(1,0) (2,0) (3,2) (6,2) (13,0) (26,4) (39,2) (78,6)
2 4 0 1	62	(1,0) (2,0) (31,0) (62,6)
3 4 0 1	31	(1,0) (31,6)
4 4 0 1	78	(1,0) (2,0) (3,0) (6,2) (13,4) (26,4) (39,4) (78,6)
0 0 1 1	50	(1,0) (2,1) (5,0) (10,5) (25,0) (50,6)
1 0 1 1	52	(1,0) (2,0) (4,2) (13,4) (26,4) (52,6)
2 0 1 1	12	(1,0) (2,0) (3,2) (4,0) (6,2) (12,6)
3 0 1 1	65	(1,1) (5,2) (13,5) (65,6)
4 0 1 1	78	(1,0) (2,0) (3,0) (6,2) (13,4) (26,4) (39,4) (78,6)
0 1 1 1	24	(1,0) (2,0) (3,2) (4,0) (6,2) (8,4) (12,2) (24,6)
1 1 1 1	60	(1,0) (2,1) (3,0) (4,3) (5,0) (6,3) (10,2) (12,5) (15,0) (20,4) (30, 4) (60,6)
2 1 1 1	130	(1,1) (2,1) (5,2) (10,2) (13,1) (26,5) (65,2) (130,6)
3 1 1 1	7	(1,0) (7,6)
4 1 1 1	62	(1,0) (2,0) (31,0) (62,6)
0 2 1 1	63	(1,0) (3,0) (7,0) (9,0) (21,0) (63,6)
1 2 1 1	20	(1,1) (2,1) (4,3) (5,4) (10,4) (20,6)
2 2 1 1	120	(1,0) (2,1) (3,0) (4,1) (5,0) (6,1) (8,1) (10,2) (12,1) (15,0) (20, 2) (24,5) (30,2) (40,2) (60,2) (120,6)
3 2 1 1	30	(1,0) (2,0) (3,2) (5,0) (6,4) (10,0) (15,4) (30,6)
4 2 1 1	31	(1,0) (31,6)
0 3 1 1	30	(1,1) (2,1) (3,1) (5,2) (6,3) (10,2) (15,2) (30,6)
1 3 1 1	60	(1,0) (2,0) (3,2) (4,2) (5,0) (6,2) (10,0) (12,4) (15,2) (20,4) (30, 2) (60,6)
2 3 1 1	42	(1,0) (2,0) (3,0) (6,0) (7,0) (14,0) (21,0) (42,6)
3 3 1 1	130	(1,0) (2,1) (5,0) (10,2) (13,0) (26,5) (65,0) (130,6)
4 3 1 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
0 4 1 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
1 4 1 1	24	(1,0) (2,0) (3,0) (4,2) (6,0) (8,2) (12,2) (24,6)
2 4 1 1	78	(1,0) (2,0) (3,0) (6,2) (13,0) (26,4) (39,0) (78,6)
3 4 1 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
4 4 1 1	30	(1,1) (2,2) (3,3) (5,2) (6,4) (10,4) (15,4) (30,6)
0 0 2 1	24	(1,0) (2,0) (3,2) (4,0) (6,2) (8,0) (12,2) (24,6)

TABLE 2. Continued

$a_0, \dots, a_k$	s	$(m, l_m(f)), m \mid s$
1 0 2 1	63	(1,0) (3,0) (7,0) (9,0) (21,0) (63,6)
2 0 2 1	60	(1,1) (2,1) (3,1) (4,3) (5,2) (6,3) (10,2) (12,5) (15,2) (20,4) (30, 4) (60,6)
3 0 2 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
4 0 2 1	60	(1,0) (2,1) (3,0) (4,1) (5,0) (6,1) (10,2) (12,5) (15,0) (20,2) (30, 2) (60,6)
0 1 2 1	40	(1,0) (2,1) (4,1) (5,0) (8,5) (10,2) (20,2) (40,6)
1 1 2 1	120	(1,1) (2,1) (3,1) (4,1) (5,2) (6,1) (8,1) (10,2) (12,1) (15,2) (20, 2) (24,5) (30,2) (40,2) (60,2) (120,6)
2 1 2 1	52	(1,0) (2,0) (4,2) (13,0) (26,4) (52,6)
3 1 2 1	39	(1,0) (3,2) (13,4) (39,6)
4 1 2 1	30	(1,0) (2,0) (3,0) (5,0) (6,2) (10,0) (15,0) (30,6)
0 2 2 1	130	(1,1) (2,1) (5,2) (10,2) (13,1) (26,5) (65,2) (130,6)
1 2 2 1	30	(1,0) (2,1) (3,2) (5,0) (6,5) (10,2) (15,2) (30,6)
2 2 2 1	52	(1,0) (2,0) (4,2) (13,4) (26,4) (52,6)
3 2 2 1	62	(1,0) (2,0) (31,0) (62,6)
4 2 2 1	63	(1,0) (3,0) (7,0) (9,0) (21,0) (63,6)
0 3 2 1	63	(1,0) (3,0) (7,0) (9,0) (21,0) (63,6)
1 3 2 1	126	(1,0) (2,0) (3,0) (6,0) (7,0) (9,0) (14,0) (18,0) (21,0) (42,0) (63, 0) (126,6)
2 3 2 1	20	(1,0) (2,1) (4,3) (5,0) (10,2) (20,6)
3 3 2 1	78	(1,0) (2,0) (3,0) (6,2) (13,4) (26,4) (39,4) (78,6)
4 3 2 1	15	(1,1) (3,3) (5,4) (15,6)
0 4 2 1	78	(1,0) (2,0) (3,0) (6,2) (13,0) (26,4) (39,0) (78,6)
1 4 2 1	62	(1,0) (2,0) (31,0) (62,6)
2 4 2 1	60	(1,0) (2,0) (3,2) (4,2) (5,0) (6,2) (10,0) (12,4) (15,4) (20,2) (30, 4) (60,6)
3 4 2 1	10	(1,1) (2,2) (5,2) (10,6)
4 4 2 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
0 0 3 1	24	(1,0) (2,0) (3,0) (4,0) (6,2) (8,0) (12,2) (24,6)
1 0 3 1	60	(1,1) (2,1) (3,1) (4,1) (5,2) (6,1) (10,2) (12,5) (15,2) (20,2) (30, 2) (60,6)
2 0 3 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
3 0 3 1	60	(1,0) (2,1) (3,2) (4,3) (5,0) (6,3) (10,2) (12,5) (15,2) (20,4) (30, 4) (60,6)
4 0 3 1	126	(1,0) (2,0) (3,0) (6,0) (7,0) (9,0) (14,0) (18,0) (21,0) (42,0) (63, 0) (126,6)
0 1 3 1	40	(1,1) (2,1) (4,1) (5,2) (8,5) (10,2) (20,2) (40,6)
1 1 3 1	15	(1,0) (3,2) (5,0) (15,6)
2 1 3 1	78	(1,0) (2,0) (3,0) (6,2) (13,0) (26,4) (39,0) (78,6)
3 1 3 1	52	(1,0) (2,0) (4,2) (13,4) (26,4) (52,6)
4 1 3 1	120	(1,0) (2,1) (3,0) (4,1) (5,0) (6,1) (8,1) (10,2) (12,1) (15,0) (20, 2) (24,5) (30,2) (40,2) (60,2) (120,6)
0 2 3 1	130	(1,0) (2,1) (5,0) (10,2) (13,4) (26,5) (65,4) (130,6)
1 2 3 1	126	(1,0) (2,0) (3,0) (6,0) (7,0) (9,0) (14,0) (18,0) (21,0) (42,0) (63, 0) (126,6)
2 2 3 1	31	(1,0) (31,6)
3 2 3 1	52	(1,0) (2,0) (4,2) (13,0) (26,4) (52,6)
4 2 3 1	30	(1,1) (2,1) (3,3) (5,2) (6,5) (10,2) (15,4) (30,6)
0 3 3 1	126	(1,0) (2,0) (3,0) (6,0) (7,0) (9,0) (14,0) (18,0) (21,0) (42,0) (63, 0) (126,6)

TABLE 2. Continued

$a_0, \dots, a_k$	s	$(m, l_m(f)), m \mid s$
1 3 3 1	30	(1,0) (2,1) (3,0) (5,0) (6,3) (10,4) (15,0) (30,6)
2 3 3 1	78	(1,0) (2,0) (3,2) (6,2) (13,0) (26,4) (39,2) (78,6)
3 3 3 1	20	(1,1) (2,1) (4,3) (5,2) (10,2) (20,6)
4 3 3 1	63	(1,0) (3,0) (7,0) (9,0) (21,0) (63,6)
0 4 3 1	39	(1,0) (3,2) (13,4) (39,6)
1 4 3 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
2 4 3 1	10	(1,1) (2,2) (5,4) (10,6)
3 4 3 1	60	(1,0) (2,0) (3,0) (4,2) (5,0) (6,2) (10,0) (12,4) (15,0) (20,2) (30, 4) (60,6)
4 4 3 1	31	(1,0) (31,6)
0 0 4 1	25	(1,1) (5,5) (25,6)
1 0 4 1	78	(1,0) (2,0) (3,2) (6,2) (13,0) (26,4) (39,2) (78,6)
2 0 4 1	130	(1,0) (2,1) (5,0) (10,2) (13,0) (26,5) (65,0) (130,6)
3 0 4 1	12	(1,0) (2,0) (3,0) (4,0) (6,2) (12,6)
4 0 4 1	52	(1,0) (2,0) (4,2) (13,0) (26,4) (52,6)
0 1 4 1	24	(1,0) (2,0) (3,0) (4,0) (6,2) (8,4) (12,2) (24,6)
1 1 4 1	31	(1,0) (31,6)
2 1 4 1	14	(1,0) (2,0) (7,0) (14,6)
3 1 4 1	130	(1,0) (2,1) (5,0) (10,2) (13,4) (26,5) (65,4) (130,6)
4 1 4 1	60	(1,1) (2,1) (3,3) (4,3) (5,2) (6,3) (10,2) (12,5) (15,4) (20,4) (30, 4) (60,6)
0 2 4 1	126	(1,0) (2,0) (3,0) (6,0) (7,0) (9,0) (14,0) (18,0) (21,0) (42,0) (63, 0) (126,6)
1 2 4 1	62	(1,0) (2,0) (31,0) (62,6)
2 2 4 1	30	(1,0) (2,0) (3,2) (5,0) (6,4) (10,0) (15,2) (30,6)
3 2 4 1	120	(1,1) (2,1) (3,1) (4,1) (5,2) (6,1) (8,1) (10,2) (12,1) (15,2) (20, 2) (24,5) (30,2) (40,2) (60,2) (120,6)
4 2 4 1	20	(1,0) (2,1) (4,3) (5,0) (10,4) (20,6)
0 3 4 1	30	(1,0) (2,1) (3,2) (5,0) (6,3) (10,2) (15,4) (30,6)
1 3 4 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
2 3 4 1	65	(1,1) (5,2) (13,5) (65,6)
3 3 4 1	21	(1,0) (3,0) (7,0) (21,6)
4 3 4 1	60	(1,0) (2,0) (3,0) (4,2) (5,0) (6,2) (10,0) (12,4) (15,0) (20,4) (30, 2) (60,6)
0 4 4 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
1 4 4 1	30	(1,1) (2,2) (3,1) (5,2) (6,4) (10,4) (15,2) (30,6)
2 4 4 1	124	(1,0) (2,0) (4,0) (31,0) (62,0) (124,6)
3 4 4 1	39	(1,0) (3,2) (13,4) (39,6)
4 4 4 1	24	(1,0) (2,0) (3,0) (4,2) (6,0) (8,2) (12,2) (24,6)